

**CHESAPEAKE BAY WATER QUALITY
MONITORING PROGRAM**

**LONG-TERM BENTHIC MONITORING
AND ASSESSMENT COMPONENT
LEVEL I COMPREHENSIVE REPORT**

JULY 1984—DECEMBER 2003 (VOLUME 1)

Prepared for

Maryland Department of Natural Resources
Resource Assessment Service
Tidewater Ecosystem Assessments
Annapolis, Maryland

Prepared by

Roberto J. Llansó
Frederick S. Kelley
Lisa C. Scott

Versar, Inc.
9200 Rumsey Road
Columbia, Maryland 21045

July 2004

FOREWORD

This document, Chesapeake Bay Water Quality Monitoring Program: Long-Term Benthic Monitoring and Assessment Component, Level I Comprehensive Report (July 1984— December 2003), was prepared by Versar, Inc., at the request of Mr. Bruce Michael of the Maryland Department of Natural Resources under Cooperative Agreement CA-03-18-07-4-30817-3734 between Versar, Inc., and the University of Maryland Center for Environmental and Estuarine Studies. The report assesses the status of Chesapeake Bay benthic communities in 2003 and evaluates their responses to changes in water quality.

ACKNOWLEDGMENTS

We are grateful to the State of Maryland's Environmental Trust Fund which partially funded this work. The benthic studies discussed in this report were conducted from the University of Maryland's research vessels and we appreciate the efforts of their captains and crew. We thank Nancy Mountford and Tim Morris of Cove Corporation who identified benthos in many of the samples and provided current taxonomic and autecological information. We also thank those at Versar whose efforts helped produce this report: the field crews who collected samples, including Martin Berlett, Donna Croson, Katherine Dillow, and Megen McBride; the laboratory technicians for processing samples, including Martin Berlett, Aaron Duprey, Bobbi Mayer, Megan McBride, Lay New, and Josh Vanderwagon; Suzanne Arcuri and Michael Winnel for taxonomic identifications; Jody Dew for managing and analyzing data; Allison Brindley for GIS support; Sherian George and Gail Lucas for document production support; and Dr. Don Strebel for web-page development.

We appreciate the efforts of Dr. Daniel M. Dauer and Anthony (Bud) Rodi of Old Dominion University who coordinate the activities of the Virginia Benthic Monitoring Program.

EXECUTIVE SUMMARY

Benthic macroinvertebrates have been an important component of the State of Maryland's Chesapeake Bay water quality monitoring program since the program's inception in 1984. Benthos integrate temporally variable environmental conditions and the effects of multiple types of environmental stress. They are sensitive indicators of environmental status. Information on the condition of the benthic community provides a direct measure of the effectiveness of management actions. This report is the 20th in a series of annual reports that summarize data up to the current sampling year. Benthic community condition and trends in the Chesapeake Bay are assessed for 2003 and compared to results from previous years.

Sampling Design and Methods

Maryland's long-term benthic monitoring program currently contains two elements: a fixed site monitoring effort directed at identifying temporal trends and a probability-based sampling effort intended to assess the areal extent of degraded benthic community condition. Benthic community condition is assessed using the benthic index of biotic integrity (B-IBI), which evaluates the ecological condition of a sample by comparing values of key benthic community attributes to reference values expected under non-degraded conditions in similar habitat types. These reference values are the benthic community restoration goals for the Chesapeake Bay. Application of the B-IBI is limited to samples collected in summer, defined as July 15 through September 30.

Twenty-seven fixed sites are sampled twice a year, in May and in late August or September. Three replicate sediment samples for benthos are collected at each fixed site with gear used since 1984. These sites are part of a more extensive suite of sites that were sampled previously at various times and locations. The probability-based sampling design is stratified simple random. It was established in 1994. Twenty-five random sites are allocated annually to each of six strata in the Maryland portion of the Chesapeake Bay. A similar stratification scheme has been used by the Commonwealth of Virginia since 1996, permitting annual estimates for the entire Chesapeake Bay. The largest portion of the Chesapeake Bay, the mainstem, is divided into three strata, and five strata consist of the major tributaries (Patuxent, Potomac, Rappahannock, York, and James rivers). Two additional strata include the remaining smaller tributaries of the Maryland upper western shore and Maryland eastern shore. The strata sampled represent the entire tidal region of the Chesapeake Bay from freshwater to polyhaline zones. Probability sites are sampled once a year in late August or September. One sample is collected at each probability site using a Young grab with a surface area of 440 cm².

All samples are sieved on a 0.5-mm screen and preserved in the field. At each site, temperature, conductivity, salinity, dissolved oxygen concentration, and pH of the water column are measured at various depths, and silt-clay percent, total organic carbon, total

inorganic carbon, and total nitrogen are measured from sediment samples processed in the laboratory.

Trends in Fixed Site Benthic Condition

Statistically significant 19-year B-IBI trends were detected at 10 of the 27 sites currently monitored. Benthic community condition declined at 4 sites and improved at 6 sites. Trends detected through 2002 were still present in 2003 with the exception of an improving trend in the Choptank River (Sta. 64) first reported in 2002, which was no longer significant with the addition of the 2003 data. New trends were detected in 2003 for Station 15 in the mainstem of the Bay (improving) and for Station 204 in the Severn River (degrading).

Sites with improving B-IBI trends were located in the main stem of the Bay (4 sites), the Elk River (Sta. 29), and the Potomac River at St. Clements Island (Sta. 51). Sites with degrading B-IBI trends were located in the Severn River (Sta. 204), the Patuxent River at Holland Cliff (Sta. 77), the Potomac River at Morgantown (Sta. 44), and the Nanticoke River (Sta. 62). Benthic organisms respond to long-term patterns in water quality parameters, such as dissolved oxygen concentrations, chlorophyll a, total nitrogen, and sediment loadings, in addition to natural fluctuations in salinity. Improving trends are likely to reflect undergoing basin-wide changes resulting from management actions. Degrading trends reflect the cumulative impacts of pollution loadings in regions with significant problems that are not yet responding to pollution abatement.

Improving trends were attributed to increases in faunal abundance at two of the mainstem sites, positive changes in the abundance of pollution-indicative organisms in the Elk River, and increases in diversity and a general improvement of the condition of the benthic community in the lower shallow Potomac River (Sta. 51) suggesting improvements in water quality in this region of the river. The Elk River trend was associated with improving trends for nutrients, chlorophyll, and sediment concentrations reported for this region of the Bay.

Degrading trends were attributed to a decrease in biomass in the Severn River, decreases in abundance and biomass in the Potomac River at Morgantown (Sta. 44), and decreases in biomass and diversity in the Nanticoke River. The upper portion of the Severn River is affected by severe hypoxia. Station 44 in the Potomac River is on the slope of the deep channel and is affected by tilts of the pycnocline bringing episodic fluctuations in dissolved oxygen and salinity. The Nanticoke River is affected by high sediment loads. Low biomass relative to reference conditions is a problem common to the Nanticoke River and the other tributaries of the lower eastern shore of Maryland.

B-IBI trends at sites in the Patuxent River gave signals of recovery. At Station 77, the magnitude of the degrading trend continued to diminish, and a long-term degrading trend at Station 71 near Broomes Island disappeared with the addition of the 2002 data.

Recovery at Station 77 was associated with increases in biomass, reflecting increasing densities of bivalves and crustaceans in the upper Patuxent River. It is not clear, however, whether these patterns reflect water quality changes or are largely influenced by changes in river flow resulting from drier than normal years since 1999.

Baywide Benthic Community Condition

The area of Chesapeake Bay estimated to fail the restoration goals in 2003 increased from 53% in 2002 to 59% in 2003, the largest estimate of degraded area since baywide monitoring began in 1996. The higher estimates for 2003 were associated with high flow conditions in the Bay, which were responsible for high nutrient and sediment run off, strong water column density stratification events, and widespread hypoxia. Wet conditions in 2003 were in contrast with below normal rainfall and low river flows in 2002. Inter-annual changes in benthic condition appear to be associated with changes in hydrology (dry versus wet years) and year-to-year fluctuations in dissolved oxygen concentrations. However, benthic community degradation in Chesapeake Bay continues to be large in any given year. In the Maryland portion of the Bay, 65% of the tidal waters failed the Chesapeake Bay benthic community restoration goals in 2003.

Fifty-one percent of the degraded Chesapeake Bay bottom in 2003 (3,501 km²) was marginally to moderately impaired. In the Maryland portion of the Bay, 37% of the degraded bottom (1,516 km²) was marginally to moderately impaired. No obvious trends in the percentage of area with marginal or moderate degradation were observed over the time series.

The Potomac, Patuxent, and York rivers were in the poorest condition among the ten bay strata in 2003. The bottom area failing the restoration goals for these three systems was 80%, 76%, and 84%, respectively. The percent severely degraded condition of the Potomac River in 2003 was the highest recorded since random monitoring began in 1994. The upper Bay mainstem and the Virginia mainstem were in best condition overall. Unlike previous years, Maryland eastern shore tributaries exhibited unusually high levels of degradation near 65 percent in 2003.

Generally, there was good agreement between the status and trends for water quality parameters and the benthic community condition. Over the period 1996-2003, high percentages of severely degraded sites failing the restoration goals due to insufficient abundance or biomass occurred in the Potomac River, Patuxent River, and the mainstem of the Chesapeake Bay. Sites with high incidence of failure due to excess abundance were most frequently located in the Maryland eastern shore tributaries, upper Bay mainstem, the James River, and the York River. Severely degraded and depauperate benthic communities are symptomatic of prolonged oxygen stress while excess abundance and biomass are symptomatic of eutrophic conditions in the absence of low dissolved oxygen stress. Low dissolved oxygen events are common and severe in the Potomac River and the Maryland

mainstem, and the Patuxent River experiences annual events of variable intensity. Maryland eastern tributaries have high agricultural land use, high nutrient input, and high chlorophyll values but low frequencies of low dissolved oxygen events. Restoration goal failure due to severely degraded benthic fauna was more common than failure due to excess abundance or biomass of benthic organisms.

TABLE OF CONTENTS

VOLUME 1

Page

FOREWORD iii

ACKNOWLEDGEMENTS v

EXECUTIVE SUMMARY vii

1.0 INTRODUCTION 1-1

 1.1 BACKGROUND 1-1

 1.2 OBJECTIVES OF THIS REPORT 1-3

 1.3 ORGANIZATION OF REPORT 1-4

2.0 METHODS 2-1

 2.1 SAMPLING DESIGN 2-1

 2.1.1 Fixed Site Sampling 2-1

 2.1.2 Probability-based Sampling 2-8

 2.2 SAMPLE COLLECTION 2-11

 2.2.1 Station Location 2-11

 2.2.2 Water Column Measurements 2-11

 2.2.3 Benthic Samples 2-14

 2.3 LABORATORY PROCESSING 2-14

 2.4 DATA ANALYSIS 2-15

 2.4.1 The B-IBI and the Chesapeake Bay Benthic Community
 Restoration Goals 2-16

 2.4.2 Fixed Site Trend Analysis 2-16

 2.4.3 Probability-Based Estimation 2-16

3.0 RESULTS 3-1

 3.1 TRENDS IN FIXED SITE BENTHIC CONDITION 3-1

 3.2 BAYWIDE BOTTOM COMMUNITY CONDITION 3-2

4.0 DISCUSSION 4-1

 4.1 PATUXENT RIVER 4-2

 4.2 POTOMAC RIVER 4-5

 4.3 UPPER WESTERN TRIBUTARIES 4-8

 4.4 EASTERN TRIBUTARIES 4-9

 4.5 MARYLAND MID BAY AND UPPER BAY MAINSTEMS 4-9

 4.6 VIRGINIA TRIBUTARIES 4-10

 4.7 CONCLUSIONS 4-10

5.0 REFERENCES 5-1

TABLE OF CONTENTS**Page****VOLUME 1****APPENDICES**

A	FIXED SITE COMMUNITY ATTRIBUTE 1985-2003 TREND ANALYSIS RESULTS.....	A-1
B	FIXED SITE B-IBI VALUES, SUMMER 2003.....	B-1
C	RANDOM SITE B-IBI VALUES, SUMMER 2003.....	C-1

VOLUME 2**DATA SUMMARIES**

A	BENTHIC ENVIRONMENT AND COMMUNITY COMPOSITION AT FIXED SITES: SPRING 2003	A-1
B	BENTHIC ENVIRONMENT AND COMMUNITY COMPOSITION AT FIXED SITES: SUMMER 2003	B-1
C	BENTHIC ENVIRONMENT AND COMMUNITY COMPOSITION AT THE MARYLAND BAY RANDOM SITES: SUMMER 2003.....	C-1

LIST OF TABLES

Table	Page
2-1 Location, habitat type, sampling gear, and habitat criteria for fixed sites	2-5
2-2 Allocation of probability-based baywide samples, 1994	2-8
2-3 Allocation of probability-based baywide samples, in and after 1995	2-11
2-4 Methods used to measure water quality parameters	2-13
2-5 Taxa for which biomass was estimated in samples collected between 1985 and 1993	2-15
3-1 Summer trends in benthic community condition, 1985-2003	3-5
3-2 Summer trends in benthic community attributes at mesohaline stations 1985- 2003	3-6
3-3 Summer trends in benthic community attributes at oligohaline and tidal freshwater stations 1985-2003	3-7
3-4 Estimated tidal area failing to meet the Chesapeake Bay benthic community restoration goals in the Chesapeake Bay, Maryland, Virginia, and each of the 10 sampling strata	3-8
3-5 Sites severely degraded and failing the restoration goals for insufficient abundance, insufficient biomass, or both as a percentage of site failing the goals, 1996 to 2003	3-11
3-6 Sites failing the restoration goals for excess abundance, excess biomass, or both as a percentage of sites failing the goals, 1996 to 2003	3-11

LIST OF FIGURES

Figure	Page
2-1 Fixed sites sampled in 2003.....	2-2
2-2 Fixed sites sampled from 1984 to 1989	2-3
2-3 Small areas and fixed sites sampled from 1989 to 1994.....	2-4
2-4 Maryland baywide sampling strata in and after 1995.....	2-9
2-5 Maryland probability-based sampling sites for 2003	2-10
2-6 Chesapeake Bay stratification scheme	2-12
3-1 Results of probability-based benthic sampling of the Maryland Chesapeake Bay and its tidal tributaries in 2003.....	3-12
3-2 Results of probability-based benthic sampling of the Virginia Chesapeake Bay and its tidal tributaries in 2003.....	3-13
3-3 Proportion of the Maryland Bay failing the Chesapeake Bay benthic community restoration goals from 1994 to 2003	3-14
3-4 Proportion of the Chesapeake Bay, Maryland, Virginia, and the 10 sampling strata failing the Chesapeake Bay benthic community restoration goals in 2003	3-15
3-5 Proportion of the Maryland sampling strata failing the Chesapeake Bay benthic community restoration goals, 1995 to 2003	3-16
3-6 Proportion of the Virginia sampling strata failing the Chesapeake Bay benthic community restoration goals, 1996 to 2003	3-17
3-7 Proportion of the Chesapeake Bay failing the Chesapeake Bay benthic community restoration goals, 1996 to 2003	3-18
4-1 Annual mean flow into Chesapeake Bay, 1937-2003	4-1
4-2 Relationship of benthic index of biotic integrity to percent dissolved oxygen observations below 2 mg/L in the mesohaline Patuxent River.....	4-3

LIST OF FIGURES (Continued)

Figure	Page
4-3 Relationship of benthic index of biotic integrity to dissolved oxygen concentration at the time of benthic sample collection in the mesohaline Patuxent River	4-3
4-4 Relationship of benthic index of biotic integrity to average chlorophyll <i>a</i> concentration in the mesohaline Patuxent River	4-4
4-5 Relationship of benthic index of biotic integrity to dissolved oxygen concentration at the time of benthic sample collection in the mesohaline Potomac River	4-5
4-6 Relationship between percent DO observations below 2 mg/L and water depth in the mesohaline Potomac River.....	4-6
4-7 Probability of observing severely degraded benthos as a function of water depth in the mesohaline Potomac River.....	4-7

1.0 INTRODUCTION

1.1 BACKGROUND

Monitoring is a necessary part of environmental management because it provides the means for assessing the effectiveness of previous management actions and the information necessary to focus future actions (NRC 1990). Towards these ends, the State of Maryland has maintained an ecological monitoring program for Chesapeake Bay since 1984. The goals of the program are to:

- quantify the types and extent of water quality problems (i.e., characterize the "state-of-the-bay");
- determine the response of key water quality measures to pollution abatement and resource management actions;
- identify processes and mechanisms controlling the bay's water quality;
- define linkages between water quality and living resources; and
- contribute information to the Water Quality Characterization Report (305b report) and the List of Impaired Waters (303d list).

The program includes elements to measure water quality, sediment quality, phytoplankton, zooplankton, and benthic macroinvertebrates (i.e., those invertebrates retained on a 0.5-mm mesh sieve). The monitoring program includes assessments of biota because the condition of biological indicators integrates temporally variable environmental conditions and the effects of multiple types of environmental stress. In addition, most environmental regulations and contaminant control measures are designed to protect biological resources; therefore, information about the condition of biological resources provides a direct measure of the effectiveness of management actions.

The Maryland program uses benthic macroinvertebrates as biological indicators because they are reliable and sensitive indicators of habitat quality in aquatic environments. Most benthic organisms have limited mobility and cannot avoid changes in environmental conditions (Gray 1979). Benthos live in bottom sediments, where exposure to contaminants and oxygen stress are most frequent. Benthic assemblages include diverse taxa representing a variety of sizes, modes of reproduction, feeding guilds, life history characteristics, and physiological tolerances to environmental conditions; therefore, they respond to and integrate natural and anthropogenic changes in environmental conditions in a variety of ways (Pearson and Rosenberg 1978; Warwick 1986; Dauer 1993; Wilson and Jeffrey 1994).

Benthic organisms are also important secondary producers, providing key linkages between primary producers and higher trophic levels (Virnstein 1977; Holland et al. 1980, 1989; Baird and Ulanowicz 1989; Diaz and Schaffner 1990). Benthic invertebrates are among the most important components of estuarine ecosystems and may represent the largest standing stock of organic carbon in estuaries (Frithsen 1989). Many benthic organisms, such as oysters and clams, are economically important. Others, such as polychaete worms and small crustaceans, contribute significantly to the diets of economically important bottom feeding juvenile and adult fishes, such as spot and croaker (Homer and Boynton 1978; Homer et al. 1980).

The Chesapeake Bay Program's decision to adopt Benthic Community Restoration Goals (Ranasinghe et al. 1994a updated by Weisberg et al. 1997) enhanced use of benthic macroinvertebrates as a monitoring tool. Based largely on data collected as part of Maryland's monitoring effort, these goals describe the characteristics of benthic assemblages expected at sites exposed to little environmental stress. The Restoration Goals provide a quantitative benchmark against which to measure the health of sampled assemblages and ultimately the Chesapeake Bay. Submerged aquatic vegetation (Dennison et al. 1993) and benthic macroinvertebrates are the only biological communities for which such quantitative goals have been established in Chesapeake Bay. Restoration goals for phytoplankton and zooplankton are under development.

A variety of anthropogenic stresses affect benthic macroinvertebrate communities in Chesapeake Bay. These include toxic contamination, organic enrichment, and low dissolved oxygen. While toxic contamination is generally restricted to urban and industrial areas typically associated with ports, low dissolved oxygen (hypoxia) is the more widespread problem encompassing an area of about 600 million m², mainly along the deep mainstem of the bay and at the mouth of the major Chesapeake Bay tributaries (Flemer et al. 1983).

A variety of factors contribute to the development and spatial variation of hypoxia in the Chesapeake Bay. Freshwater inflow, salinity, temperature, wind stress, and tidal circulation are primary factors in the development of hypoxia (Holland et al. 1987; Tuttle et al. 1987; Boicourt 1992). The development of vertical salinity gradients during the spring freshwater run off leads to water column density stratification. The establishment of a pycnocline, in association with periods of calm and warm weather, restricts water exchange between the surface and the bottom layers of the estuary, where oxygen consumption is large. The formation or the disruption of the pycnocline is probably the most important process determining the intensity and extent of hypoxia (Seliger et al. 1985; Boicourt 1992), albeit not the only one. Biological processes contribute significantly to deep water oxygen depletion in Chesapeake Bay (Officer et al. 1984). Benthic metabolic rates increase during spring and early summer, leading to an increase of the rate of oxygen consumption in bottom waters. This depends in part on the amount of organic carbon available for the benthos, which is derived to a large extent from seasonal phytoplankton blooms (Officer et al. 1984). Anthropogenic nutrient inputs to the Chesapeake Bay further stimulate phytoplankton growth, which results in increased deposition of

organic matter to the sediments and a concomitant increase in chemical and biological oxygen demand (Malone 1987). Winter to spring accumulation of phytoplankton biomass has been linked to depletion of bottom water oxygen in Chesapeake Bay (Malone et al. 1988; Boynton and Kemp 2000).

The effects of hypoxia on benthic organisms vary as a function of the severity, spatial extent, and duration of the low dissolved oxygen event. Oxygen concentrations down to about 2 mg l⁻¹ do not appear to significantly affect benthic organisms, although incipient community effects have been measured at 3 mg l⁻¹ (Diaz and Rosenberg 1995; Ritter and Montagna 1999). Hypoxia brings about structural and organizational changes in the community, and may lead to hypoxia resistant communities. With an increase in the frequency of hypoxic events, benthic populations become dominated by fewer and short-lived species, and their overall productivity is decreased (Diaz and Rosenberg 1995). Major reductions in species number and abundance in the Chesapeake Bay have been attributed to hypoxia (Llansó 1992). These reductions become larger both spatially and temporally as the severity and duration of hypoxic events increase. As hypoxia becomes persistent, mass mortality of benthic organisms often occurs with almost complete elimination of the macrofauna.

Hypoxia has also major impacts on the survival and behavior of a variety of benthic organisms and their predators (Diaz and Rosenberg 1995). Many infaunal species respond to low oxygen by migrating toward the sediment surface, thus potentially increasing their availability to demersal predators. On the other hand, reduction or elimination of the benthos following severe hypoxic or anoxic (no oxygen) events may result in a reduction of food for demersal fish species and crabs. Therefore, the structural changes and species replacements that occur in communities affected by hypoxia may alter the food supply of important ecological and economical fish species in Chesapeake Bay. Given that dissolved oxygen and nutrient inputs are critical factors in the health of the resources of the Chesapeake Bay region, monitoring that evaluates benthic community condition and tracks changes over time helps Chesapeake Bay managers assess the effectiveness of nutrient reduction efforts and the status of the biological resources of one of the largest and most productive estuaries in the nation.

1.2 OBJECTIVES OF THIS REPORT

This report marks the 20th in a series of Level I Comprehensive reports produced annually by the Long-Term Benthic Monitoring and Assessment Component (LTB) of the Maryland Chesapeake Bay Water Quality Monitoring Program. Level I reports summarize data from the latest sampling year and provide a limited examination of how conditions in the latest year differ from conditions in previous years of the study, as well as how data from this year contribute to describing trends in the bay's condition.

The report reflects the maturity of the current program's focus and design. Approaches introduced when the new program design was implemented in 1995 continue

to be extended, developed, and better defined. The level of detail in which changes are examined at the fixed stations sampled for trend analysis continues to increase. For example, we report on how species contribute to changes in condition and discuss results in relation to changes in water quality. The Benthic Index of Biotic Integrity (B-IBI) is applied to each sampling site, from tidal freshwater to polyhaline habitats, and thus provides an uniform measure of ecological condition across the estuarine gradient. In describing baywide benthic community condition, estimates of degraded condition are presented for at least eight years for all subregions of the Bay, and community measures that contribute to Restoration Goal failure are used to diagnose the causes of failure.

The continued presentation of estimates of Bay area meeting the Chesapeake Bay Program's Benthic Community Restoration Goals, rather than Maryland estimates only, reflects improved coordination and unification of objectives among the Maryland and Virginia benthic monitoring programs. The sampling design and methods in both states are compatible and complementary.

In addition to the improvements in technical content, we have enhanced electronic production and transmittal of data. This report is produced in Adobe Acrobat format to facilitate distribution across the internet. Data and program information are available to the research community and the general public through the Chesapeake Bay Benthic Monitoring Program Home Page on the World-Wide-Web at <http://www.baybenthos.versar.com>. Expansion of the site continues, with new program information, data, and documents being added every year. The 2003 data, as well as the data from previous years, can be downloaded from this site. The Benthic Monitoring Program Home Page represents the culmination of collaborative efforts between Versar, Maryland DNR, and the Chesapeake Information Management System (CIMS). The activities that Versar undertakes as a partner of CIMS were recorded in a Memorandum of Agreement signed October 28, 1999.

1.3 ORGANIZATION OF REPORT

This report has two volumes. Volume 1 is organized into four major sections and three appendices. Section 1 is this introduction. Section 2 presents the field, laboratory, and data analysis methods used to collect, process, and evaluate LTB samples. Section 3 presents the results of analyses conducted for 2003 and previous years, and consists of two assessments: an assessment of trends in benthic community condition at sites sampled annually by LTB in the Maryland Chesapeake Bay, and an assessment of the area of the Bay that meets the Chesapeake Bay Benthic Community Restoration Goals. Section 4 discusses the results and evaluates status and trends relative to recent changes in water quality. Section 5 is the literature cited in the report. Appendix A amplifies information presented in Table 3-2 by providing p-values and rates of change for the 1985-2003 fixed site trend analysis. Finally, Appendices B and C present the B-IBI values for the 2003 fixed and random samples, respectively. Volume 2 consists of the benthic, sedimentary, and hydrographic data appendices.

2.0 METHODS

2.1 SAMPLING DESIGN

The LTB sampling program contains two primary elements: a fixed site monitoring effort directed at identifying trends in benthic condition and a probability-based sampling effort intended to estimate the area of the Maryland Chesapeake Bay with benthic communities meeting the Chesapeake Bay Program's benthic community restoration goals (Ranasinghe et al. 1994a, updated by Weisberg et al. 1997; Alden et al. 2002). The sampling design for each of these elements is described below.

2.1.1 Fixed Site Sampling

The fixed site element of the program involves sampling at 27 sites, 23 of which have been sampled since the program's inception in 1984, 2 since 1989, and 2 since 1995 (Figure 2-1). Sites are defined by geography (within 1 km from a fixed location), and by specific depth and substrate criteria (Table 2-1).

The 2003 fixed site sampling continues trend measurements, which began with the program's initiation in 1984. In the first five years of the program, from July 1984 to June 1989, 70 fixed stations were sampled 8 to 10 times per year. On each visit, three benthic samples were collected at each site and processed. Locations of the 70 fixed sites are shown in Figure 2-2.

In the second five years of the program, from July 1989 to June 1994, fixed site sampling was continued at 29 sites and a stratified random sampling element was added. Samples were collected at random from approximately 25 km² small areas surrounding these sites (Figure 2-3) to assess the representativeness of the fixed locations. Sites 06, 47, 62, and 77, which are part of the current design, were not sampled during this five-year period. Stratum boundaries were delineated on the basis of environmental factors that are important in controlling benthic community distributions: salinity regime, sediment type, and bottom depth (Holland et al. 1989). In addition, four new areas were established in regions of the Bay targeted for management actions to abate pollution: the Patuxent River, Choptank River, and two areas in Baltimore Harbor. Each area was sampled four to six times each year.

From July 1994 to the present, three replicate samples were collected in spring and summer at most of the current suite of 27 sites (Stations 203 and 204 were added in 1995, Table 2-1, Fig. 2-1). This sampling regime was selected as being most cost effective after analysis of the first 10 years of data jointly with the Virginia Benthic Monitoring Program (Alden et al. 1997).




Figure 2-1. Fixed sites sampled in 2003




Figure 2-2. Fixed sites sampled from 1984 to 1989; some of these sites are part of the current design




Figure 2-3. Small areas and fixed sites sampled from 1989 to 1994

Table 2-1. Location, habitat type, (Table 5, Weisberg et al. 1997), sampling gear, and habitat criteria for fixed sites									
Stratum	Sub-Estuary	Habitat	Station	Latitude (NAD 83)	Longitude (NAD 83)	Sampling Gear	Habitat Criteria		
							Depth (m)	Siltclay (%)	Distance (km)
Potomac River	Potomac River	Tidal Freshwater	036	38.769781	77.037531	WildCo Box Corer	< = 5	> = 40	1.0
		Oligohaline	040	38.357458	77.230534	WildCo Box Corer	6.5-10	> = 80	1.0
		Low Mesohaline	043	38.384125	76.989028	Modified Box Corer	< = 5	< = 30	1.0
		Low Mesohaline	047	38.365125	76.984695	Modified Box Corer	< = 5	< = 30	0.5
		Low Mesohaline	044	38.385625	76.995695	WildCo Box Corer	11-17	> = 75	1.0
		High Mesohaline Sand	051	38.205462	76.738020	Modified Box Corer	< = 5	< = 20	1.0
		High Mesohaline Mud	052	38.192297	76.747687	WildCo Box Corer	9-13	> = 60	1.0
Patuxent River	Patuxent River	Tidal Freshwater	079	38.750448	76.689020	WildCo Box Corer	< = 6	> = 50	1.0
		Low Mesohaline	077	38.604452	76.675017	WildCo Box Corer	< = 5	> = 50	1.0
		Low Mesohaline	074	38.547288	76.674851	WildCo Box Corer	< = 5	> = 50	0.5
		High Mesohaline Mud	071	38.395124	76.548844	WildCo Box Corer	12-18	> = 70	1.0

Table 2-1. (Continued)									
Stratum	Sub-Estuary	Habitat	Station	Latitude (NAD 83)	Longitude (NAD 83)	Sampling Gear	Habitat Criteria		
							Depth (m)	Siltclay (%)	Distance (km)
Upper Western Tributaries	Patapsco River	Low Mesohaline	023	39.208275	76.523352	WildCo Box Corer	4-7	> = 50	1.0
	Middle Branch	Low Mesohaline	022	39.254940	76.587354	WildCo Box Corer	2-6	> = 40	1.0
	Bear Creek	Low Mesohaline	201	39.234275	76.497184	WildCo Box Corer	2-4.5	> = 70	1.0
	Curtis Bay	Low Mesohaline	202	39.217940	76.563853	WildCo Box Corer	5-8	> = 60	1.0
	Back River	Oligohaline	203	39.275107	76.446015	Young-Grab	1.5-2.5	> = 80	1.0
	Severn River	High Mesohaline Mud	204	39.006778	76.504683	Young-Grab	5-7.5	> = 50	1.0
Eastern Tributaries	Chester River	Low Mesohaline	068	39.132941	76.078679	WildCo Box Corer	4-8	> = 70	1.0
	Choptank River	Oligohaline	066	38.801447	75.921825	WildCo Box Corer	< = 5	> = 60	1.0
		High Mesohaline Mud	064	38.590464	76.069340	WildCo Box Corer	7-11	> = 70	1.0
	Nanticoke River	Low Mesohaline	062	38.383952	75.849988	Petite Ponar Grab	5-8	> = 75	1.0

Table 2-1. (Continued)									
Stratum	Sub-Estuary	Habitat	Station	Latitude (NAD 83)	Longitude (NAD 83)	Sampling Gear	Habitat Criteria		
							Depth (m)	Siltclay (%)	Distance (km)
Upper Bay	Elk River	Oligohaline	029	39.479615	75.944499	WildCo Box Corer	3-7	> = 40	1.0
	Mainstem	Low Mesohaline	026	39.271441	76.290011	WildCo Box Corer	2-5	> = 70	1.0
		High Mesohaline Mud	024	39.122110	76.355346	WildCo Box Corer	5-8	> = 80	1.0
Mid Bay	Mainstem	High Mesohaline Sand	015	38.715118	76.513677	Modified Box Corer	< = 5	< = 10	1.0
		High Mesohaline Sand	001	38.419956	76.416672	Modified Box Corer	< = 5	< = 20	1.0
		High Mesohaline Sand	006	38.442456	76.443006	Modified Box Corer	< = 5	< = 20	0.5

2.1.2 Probability-based Sampling

The second sampling element, which was instituted in 1994, was probability-based summer sampling designed to estimate the area of the Maryland Chesapeake Bay and its tributaries that meet the Chesapeake Bay benthic community restoration goals (Ranasinghe et al. 1994a, updated by Weisberg et al. 1997; Alden et al. 2002). Different probability sample allocation strategies were used in 1994 than in later years. In 1994, the design was intended to estimate impaired area for the Maryland Bay and one sub-region, while in later years the design targeted five additional sub-regions as well.

The 1994 sample allocation scheme was designed to produce estimates for the Maryland Bay and the Potomac River. The Maryland Bay was divided into three strata with samples allocated unequally among them (Table 2-2); sampling intensity in the Potomac was increased to permit estimation of degraded area with adequate confidence, while mainstem and other tributary and embayment samples were allocated in proportion to their area.

Stratum	Area		Number of Samples
	km ²	%	
Maryland Mainstem (including Tangier and Pocomoke Sounds)	3,611	55.5	27
Potomac River	1,850	28.4	28
Other tributaries and embayments	1,050	16.1	11

In subsequent years, the stratification scheme was designed to produce an annual estimate for the Maryland Bay and six subdivisions. Samples were allocated equally among strata (Figure 2-4, Table 2-3). According to this allocation, a fresh new set of sampling sites were selected each year. Figure 2-5 shows the locations of the probability-based Maryland sampling sites for 2003. Regions of the Maryland mainstem deeper than 12 m were not included in sampling strata because these areas are subjected to summer anoxia and have consistently been found to be azoic.

A similar stratification scheme has been used by the Commonwealth of Virginia since 1996, permitting annual estimates for the extent of area meeting the benthic community restoration goals for the entire Chesapeake Bay (Table 2-3, Figure 2-6). These samples were collected and processed, and the data analyzed by the Virginia Chesapeake Bay Benthic Monitoring Program.




Figure 2-4. Maryland baywide sampling strata in and after 1995




Figure 2-5. Maryland probability-based sampling sites for 2003

Table 2-3. Allocation of probability-based baywide samples, in and after 1995. Maryland areas exclude 676 km² of mainstem habitat deeper than 12 m. Virginia strata were sampled by the Virginia Chesapeake Bay Benthic Monitoring Program commencing in 1996.

State	Stratum	Area			Number of Samples
		km2	State %	Bay %	
Maryland	Deep Mainstem	676	10.8	5.8	0
	Mid Bay Mainstem	2,552	40.9	22.0	25
	Eastern Tributaries	534	8.6	4.6	25
	Western Tributaries	292	4.7	2.5	25
	Upper Bay Mainstem	785	12.6	6.8	25
	Patuxent River	128	2.0	1.1	25
	Potomac River	1,276	20.4	11.0	25
	TOTAL	6,243	100.0	53.8	150
Virginia	Mainstem	4,120	76.8	35.5	25
	Rappahannock River	372	6.9	3.2	25
	York River	187	3.5	1.6	25
	James River	684	12.8	5.9	25
	TOTAL	5,363	100.0	46.2	100

2.2 SAMPLE COLLECTION

2.2.1 Station Location

From July 1984 to June 1996, stations were located using Loran-C. After June 1996 stations were located using a differential Global Positioning System. The NAD83 coordinate system is currently used.

2.2.2 Water Column Measurements

Water column vertical profiles of temperature, conductivity, salinity, dissolved oxygen concentration (DO), and pH were measured at each site. Oxidation reduction potential (ORP) was measured prior to 1996. For fixed sites, profiles consisted of water quality measurements at 1 m intervals from surface to bottom at sites 7 m deep or less, and at 3 m intervals, with additional measurements at 1.5 m intervals in the vicinity of the pycnocline, at sites deeper than 7 m. Surface and bottom measurements were made at all other sampling sites. Table 2-4 lists the measurement methods used.




Figure 2-6. Chesapeake Bay stratification scheme

Table 2-4. Methods used to measure water quality parameters.		
Parameter	Period	Method
Temperature	July 1984 to November 1984	Thermistor attached to Beckman Model RS5-3 salinometer
	December 1984 to December 1995	Thermistor attached to Hydrolab Surveyor II
	January 1996 to present	Thermistor attached to Hydrolab DataSonde 3 or YSI-6600 Sonde
Salinity and Conductivity	July to November 1984	Beckman Model RS5-3 salinometer toroidal conductivity cell with thermistor temperature compensation
	December 1984 to December 1995	Hydrolab Surveyor II nickel six-pin electrode-salt water cell block combination with automatic temperature compensation
	January 1996 to present	Hydrolab DataSonde 3 or YSI-6600 Sonde nickel six-pin electrode-salt water cell block combination with automatic temperature compensation
Dissolved Oxygen	July to November 1984	YSI Model 57 or Model 58 Oxygen Meter with automatic temperature and manual salinity compensation
	December 1984 to December 1995	Hydrolab Surveyor II membrane design probe with automatic temperature and salinity compensation
	January 1996 to present	Hydrolab DataSonde 3 or YSI-6600 Sonde membrane design probe with automatic temperature and salinity compensation
pH	July to November 1984	Orion analog pH meter with Ross glass combination electrode manually compensated for temperature
	December 1984 to December 1995	Hydrolab Surveyor II glass pH electrode and Lazaran reference electrode automatically compensated for temperature
	January 1996 to present	Hydrolab DataSonde 3 or YSI-6600 Sonde glass pH electrode and standard reference (STDREF) electrode automatically compensated for temperature
Oxidation Reduction Potential	December 1984 to December 1995	Hydrolab Surveyor II platinum banded glass ORP electrode

2.2.3 Benthic Samples

Samples were collected using four kinds of gear depending on the program element and habitat type. For the fixed site element (Table 2-1), a hand-operated box corer ("modified box corer"), which samples a 250 cm² area to a depth of 25 cm, was used in the nearshore shallow sandy habitats of the mainstem bay and tributaries. A Wildco box corer, which samples an area of 225 cm² to a depth of 23 cm, was used in shallow muddy or deep-water (> 5 m) habitats in the mainstem bay and tributaries. A Petite Ponar Grab, which samples 250 cm² to a depth of 7 cm, was used at the fixed site in the Nanticoke River to be consistent with previous sampling in the 1980s. At the two fixed sites first sampled in 1995 and at all probability-based sampling sites, a Young Grab, which samples an area of 440 cm² to a depth of 10 cm, was used.

Sample volume and penetration depth were measured for all samples; Wildco and hand-operated box cores penetrating less than 15 cm, and Young and Petite Ponar grabs penetrating less than 7 cm into the sediment were rejected and the site was re-sampled.

In the field, samples were sieved through a 0.5-mm screen using an elutriative process. Organisms and detritus retained on the screen were transferred into labeled jars and preserved in a 10% formaldehyde solution stained with rose bengal (a vital stain that aids in separating organisms from sediments and detritus).

Two surface-sediment sub-samples of approximately 120 ml each were collected for grain-size, carbon, and nitrogen analysis from an additional grab sample at each site. Surface sediment samples were frozen until they were processed in the laboratory.

2.3 LABORATORY PROCESSING

Organisms were sorted from detritus under dissecting microscopes, identified to the lowest practical taxonomic level (most often species), and counted. Oligochaetes and chironomids were mounted on slides and examined under a compound microscope for genus and species identification.

Ash-free dry weight biomass was determined by three comparable techniques during the sampling period. For samples collected from July 1984 to June 1985, biomass was directly measured using an analytical balance for major organism groups (e.g., polychaetes, molluscs, and crustaceans). Ash-free dry weight biomass was determined by drying the organisms to a constant weight at 60 EC and ashing in a muffle furnace at 500 EC for four hours. For samples collected between July 1985 and August 1993, a regression relationship between ash-free dry weight biomass and size of morphometric characters was defined for 22 species (Ranasinghe et al. 1993). The biomass of the 22 selected species was estimated from these regression relationships. These taxa (Table 2-5) were selected because they accounted for more than 85% of the abundance (Holland et al. 1988). After August 1993, ash-free dry weight biomass was measured directly for

each species by drying the organisms to a constant weight at 60 EC and ashing in a muffle furnace at 500 EC for four hours.

Table 2-5. Taxa for which biomass was estimated in samples collected between 1985 and 1993.	
Polychaeta	Mollusca
<i>Eteone heteropoda</i>	<i>Acteocina canaliculata</i>
<i>Glycinde solitaria</i>	<i>Corbicula fluminea</i>
<i>Heteromastus filiformis</i>	<i>Gemma gemma</i>
<i>Marenzelleria viridis</i>	<i>Haminoe solitaria</i>
<i>Neanthes succinea</i>	<i>Macoma balthica</i>
<i>Paraprionospio pinnata</i>	<i>Macoma mitchelli</i>
<i>Streblospio benedicti</i>	<i>Mulinia lateralis</i>
	<i>Mya arenaria</i>
	<i>Rangia cuneata</i>
	<i>Tagelus plebeius</i>
Crustacea	
<i>Cyathura polita</i>	
<i>Gammarus</i> spp.	
<i>Leptocheirus plumulosus</i>	
Miscellaneous	
<i>Carinoma tremaphoros</i>	
<i>Micrura leidyi</i>	

Silt-clay composition and carbon and nitrogen content were determined for one of the two sediment sub-samples collected at each sampling site. The other sample was archived for quality assurance purposes (Scott et al. 1988). Sand and silt-clay particles were separated by wet-sieving through a 63- μ m, stainless steel sieve and weighed using the procedures described in the Versar, Inc., Laboratory Standard Operating Procedures (Versar 1999). Carbon and nitrogen content of dried sediments was determined using an elemental analyzer; sediment carbon content was measured with a Perkin-Elmer Model 240B analyzer from 1984 to 1988, and an Exeter Analytical Inc., Model CE-440 analyzer in and after 1995. The results from both instruments are comparable. Samples are combusted at high temperature (975 °C) and the carbon dioxide and nitrogen produced are measured by thermal conductivity detection. Prior to combustion, each sample is homogenized and oven-dried.

2.4 DATA ANALYSIS

Analyses for the fixed site and probability-based elements of LTB were both performed in the context of the Chesapeake Bay Program's benthic community restoration goals and the Benthic Index of Biotic Integrity (B-IBI) by which goal attainment is

measured. The B-IBI, the Chesapeake Bay benthic community restoration goals, and statistical analysis methods for the two LTB elements are described below.

2.4.1 The B-IBI and the Chesapeake Bay Benthic Community Restoration Goals

The B-IBI is a multiple-attribute index developed to identify the degree to which a benthic assemblage meets the Chesapeake Bay Program's benthic community restoration goals (Ranasinghe et al. 1994a, updated by Weisberg et al. 1997; Alden et al. 2002). The B-IBI provides a means for comparing relative condition of benthic invertebrate assemblages across habitat types. It also provides a validated mechanism for integrating several benthic community attributes indicative of habitat "health" into a single number that measures overall benthic community condition.

The B-IBI is scaled from 1 to 5, and sites with values of 3 or more are considered to meet the restoration goals. The index is calculated by scoring each of several attributes as either 5, 3, or 1 depending on whether the value of the attribute at a site approximates, deviates slightly from, or deviates strongly from values found at the best reference sites in similar habitats, and then averaging these scores across attributes. The criteria for assigning these scores are numeric and depend on habitat. Data from seasons for which the B-IBI has not been developed were not used for B-IBI based assessment.

Benthic community condition was classified into four levels based on the B-IBI. Values less than or equal to 2 were classified as severely degraded; values from 2 to 2.6 were classified as degraded; values greater than 2.6 but less than 3.0 were classified as marginal; and values of 3.0 or more were classified as meeting the goals. Values in the marginal category do not meet the restoration goals, but they differ from the goals within the range of measurement error typically recorded between replicate samples.

2.4.2 Fixed Site Trend Analysis

Trends in condition at the fixed sites were identified using the nonparametric technique of van Belle and Hughes (1984). This procedure is based on the Mann-Kendall statistic and consists of a sign test comparing each value with all values measured in subsequent periods. The ratio of the Mann-Kendall statistic to its variance provides a normal deviate that is tested for significance. Alpha was set to 0.1 for these tests because of the low power for trend detection for biological data. An estimate of the magnitude of each significant trend was obtained using Sen's (1968) procedure which is closely related to the Mann-Kendall test. Sen's procedure identifies the median slope among all slopes between each value and all values measured in subsequent periods.

2.4.3 Probability-Based Estimation

The Maryland Bay was divided into three strata (Bay Mainstem, Potomac River, other tributaries and embayments) in 1994 (Table 2-2). It was divided into six strata in

and after 1995 (Figure 2-4, Table 2-3). The Virginia Bay was divided into four strata, beginning in 1996 (Figure 2-6, Table 2-3).

To estimate the amount of area in the entire Bay that failed to meet the Chesapeake Bay benthic community restoration goals (P), we defined for every site i in stratum h a variable y_{hi} that had a value of 1 if the benthic community met the goals, and 0 otherwise. For each stratum, the estimated proportion of area meeting the goals, p_h , and its variance were calculated as the mean of the y_{hi} 's and its variance, as follows:

$$p_h = \bar{y}_h = \sum_{i=1}^{n_h} \frac{y_{hi}}{n_h} \quad (1)$$

and

$$\text{var}(p_h) = s_h^2 = \sum_{i=1}^{n_h} \frac{(y_{hi} - \bar{y}_h)^2}{n_h - 1} \quad (2)$$

Estimates for strata were combined to achieve a statewide estimate as:

$$\hat{P}_{ps} = \bar{y}_{ps} = \sum_{h=1}^6 W_h \bar{y}_h \quad (3)$$

where the weighting factor $W_h = A_h/A$; A_h is the total area of the h th stratum, and A is the combined area of all strata. The variance of (3) was estimated as:

$$\text{var}(\hat{P}_{ps}) = \text{var}(\bar{y}_{ps}) = \sum_{h=1}^6 W_h^2 s_h^2 / n_h \quad (4)$$

The standard error for individual strata is estimated as the square root of (2), and for the combined strata, as the square root of (4).

3.0 RESULTS

3.1 TRENDS IN FIXED SITE BENTHIC CONDITION

Trend analysis is conducted on 27 fixed sites located throughout the Bay and its tributaries to assess whether benthic community condition is changing. The sites are sampled yearly in the spring and summer but the trend analysis is performed on the summer data only in order to apply the B-IBI (Weisberg et al. 1997; Alden et al. 2002). B-IBI calculations and trend analysis methods are described in Section 2.4.

The B-IBI is the primary measure used in trend analysis because it integrates several benthic community attributes into a measure of overall condition. It provides context for interpretation of observed trends because status has been calibrated to reference conditions. Significant trends that result in a change of status (sites that previously met the Chesapeake Bay benthic community restoration goals which now fail, or vice versa) are of greater management interest than trends which do not result in a change. As a first step in identifying causes of changes in condition, trends on individual attributes are identified and examined.

Table 3-1 presents trends in benthic community condition from 1985 to the present. Although the Maryland benthic monitoring component began sampling in 1984, data collected in the first year of our program were excluded from analysis to facilitate comparison of results with other components of the monitoring program. Several components of the Maryland program as well as the Virginia benthic monitoring program did not start sampling until 1985. Nineteen-year (1985-2003) trends are presented for 23 of the 27 trend sites, 15-year trends are presented for two sites in Baltimore Harbor (Stations 201 and 202) first sampled in 1989, and 9-year trends are presented for two western shore tributaries (Back River, Station 203; and Severn River, Station 204) first sampled in 1995. Trend site locations are shown in Figure 2-1.

Statistically significant B-IBI trends ($p < 0.1$) were detected at 10 of the 27 sites (Table 3-1). Benthic community condition declined at 4 sites (significantly decreasing B-IBI trend) and improved at 6 sites. Currently, 11 stations meet the goals and 16 fail the goals. Initially, 10 stations met the goals and 17 failed the goals (Table 3-1). Three stations with a significant trend have changed status relative to the first three years of the monitoring program. Stations 01 and 06 (mainstem) have improved from initial failure to currently meeting the goals (Table 3-1). Station 204 has declined in status from initially meeting the goals to currently failing the goals (Table 3-1). Station 44 (Potomac River, Morgantown) has declined from a marginal to a severely degraded condition. Last year, we reported improvements in the status of two additional stations with significant trends, from initial failure to meeting the goals. One of these two stations (Station 29, Elk River) still exhibits a positive B-IBI trend, but the status in 2003 has declined to marginal. Last year, we also reported a decline in the status of Station 62 (Nanticoke River). While this station continues to exhibit a degrading B-IBI trend, its status is improving and met the goals in 2003. Changes in status relative to those reported last year are highlighted in Table 3-1.

Significant trends present with the analysis of 2002 data were still present with the addition of the 2003 data except for Stations 15 (mainstem, North Beach), 64 (Choptank River), and 204 (Severn River). New trends are reported this year for North Beach (improving) and the Severn River (degrading). The positive Choptank River trend first observed in 2002 disappeared with the addition of the 2003 data.

Trends in community attributes that are components of the B-IBI are presented in Table 3-2 (mesohaline stations), Table 3-3 (oligohaline and tidal freshwater stations), and Appendix A.

3.2 BAYWIDE BOTTOM COMMUNITY CONDITION

The fixed site monitoring provides useful information about trends in the condition of benthic biological resources at 27 locations in the Maryland Bay but it does not provide an integrated assessment of the Bay's overall condition. The fixed sites were selected for trend monitoring because they are located in areas subject to management action and, therefore, are likely to undergo change. Because these sites were selected subjectively, there is no objective way of weighting them to obtain an unbiased estimate of Maryland baywide status.

An alternative approach for quantifying status of the bay, which was first adopted in the 1994 sampling program, is to use probability-based sampling to estimate the bottom area populated by benthos meeting the Chesapeake Bay benthic community restoration goals. Where the fixed site approach quantifies change at selected locations, the probability sampling approach quantifies the spatial extent of problems. While both approaches are valuable, developing and assessing the effectiveness of a Chesapeake Bay management strategy requires understanding the extent and distribution of problems throughout the Bay, instead of only assessing site-specific problems. Our probability-based sampling element is intended to provide that information, as well as a more widespread baseline data set for assessing the effects of unanticipated future contamination (e.g., oil or hazardous waste spills). Probability-based sampling information is also used for Chesapeake Bay aquatic life use support decisions under the Clean Water Act.

Probability-based sampling has been employed previously by LTB, but the sampled area included only 16% of the Maryland Bay (Ranasinghe et al. 1994a) which was insufficient to characterize the entire Bay. Probability-based sampling was also used in the Maryland Bay by the U.S. EPA Environmental Monitoring and Assessment Program (EMAP), but at a sampling density too low to develop precise condition estimates for the Maryland Bay. The 2003 sampling continues with efforts initiated in 1994 to develop area-based bottom condition statements for the Maryland Bay.

Estimates of tidal bottom area meeting the benthic community restoration goals are also included for the entire Chesapeake Bay. The estimates were enabled by including a probability-based sampling element in the Virginia Benthic Monitoring Program starting in 1996. The Virginia sampling is compatible and complementary to the Maryland effort and

is part of a joint effort by the two programs to assess the extent of “healthy” tidal bottom baywide.

This section presents the results of the 2003 Maryland and Virginia tidal waters probability-based sampling and adds a tenth year of results to LTB’s Maryland Bay time series. The analytical methods for estimating the areal extent of bay bottom meeting the restoration goals were presented in Section 2.0. The physical data associated with the benthic samples (bottom water salinity, temperature, DO, and sediment silt-clay and organic carbon content) can be found in the appendices (Volume 2). Only summer data (July 15-September 30) are used for the probability-based assessments.

Of the 150 Maryland samples collected with the probability-based design in 2003, 55 met and 95 failed the Chesapeake Bay benthic community restoration goals (Figure 3-1). Of the 250 probability samples collected in the entire Chesapeake Bay in 2003, 93 met and 157 failed the restoration goals. The Virginia sampling results are presented in Figure 3-2. In terms of number of sites failing the goals in Chesapeake Bay, 2003 was the worst year since probability-based sampling started 1994.

The percent degraded area in the Maryland Bay in 2003 did not change relative to the 2002 estimate, but the area with marginal benthic condition decreased substantially (Figure 3-3). The magnitude of the severely degraded condition has not changed appreciably since 1994. Results from the individual sites were weighted based on the area of the stratum represented by the site in the stratified sampling design to estimate the tidal Maryland area failing the restoration goals. In both 2002 and 2003, 65% ($\pm 5\%$ SE) of the Maryland Bay was estimated to fail the restoration goals. Expressed as area, $4,086 \pm 188 \text{ km}^2$ of the tidal Maryland Chesapeake Bay remained to be restored.

In 2003, the Potomac and the Patuxent Rivers were in the poorest condition among the six Maryland strata (Figure 3-4). The percent severely degraded condition of the Potomac River in 2003 was the highest recorded since random monitoring there began in 1994 (Figure 3-5, Table 3-4). The level of degradation in the Maryland mid-Bay mainstem and the upper western shore tributaries continued to be high in 2003, and the combined ‘severely degraded’ and ‘degraded’ categories for each of these strata were higher in 2003 than in the previous year. In 2003, the upper Bay mainstem and the eastern shore tributaries also exhibited unusual high levels of degradation (Figure 3-5). Generally, these two last strata have good benthic community condition relative to the other bay strata. Over the ten-year time series (1994-2003), more than half of the tidal Potomac River ($714-1,173 \text{ km}^2$) failed the restoration goals each year (Figure 3-5) and a large portion of that area, ranging from 48-93% ($510-867 \text{ km}^2$, Table 3-4), was severely degraded. The Maryland mid-Bay mainstem continued to have the largest amount of degraded area among the strata: $2,106 \text{ km}^2$ in 2003 (Table 3-4). Sixty-one percent of this area (including the deep trough) was severely degraded. Least severely degraded among the strata usually is the Maryland eastern shore, but again 2003 was an exception (Table 3-4).

The area of Chesapeake Bay estimated to fail the restoration goals in 2003 was the highest of the time series (Figure 3-7), but the change over the previous year can be

considered within the uncertainty margin of the estimate. Weighting results from the 250 probability sites in Maryland and Virginia, 59% ($\pm 4\%$) or $6,852 \pm 307$ km² of the tidal Chesapeake Bay was estimated to fail the restoration goals in 2003 (Table 3-4). The percentage for previous years ranged from 45% ($\pm 4\%$) in 1996 to 58% ($\pm 4\%$) in 1998 (Table 3-4). About 25% of the Chesapeake Bay continued to exhibit severely degraded benthic condition.

In Virginia, levels of degradation for all tributaries in 2003 were high and in tune with the pronounced degradation seen bay-wide (Figure 3-4, Table 3-4). Benthic community condition in all Virginia tributaries declined in 2003 relative to the previous year, with the York River exhibiting the largest decline and the more extensive degradation (Figure 3-6). Eighty-four percent of the York River had degraded benthic community in 2003. The Virginia mainstem, which usually supports good benthos, exhibited the largest increase in area degraded of the eight-year time series, almost reaching 50% degradation.

As reported in previous years, and for the period 1996-2003, five strata (Potomac River, Patuxent River, mid-Bay mainstem, upper western tributaries, and Virginia mainstem) had a large percentage (>65%) of sites failing the goals because of insufficient abundance or biomass of organisms relative to reference conditions (Table 3-5). Except for the Virginia mainstem, these strata also had a high percentage (>55%) of failing sites classified as severely degraded (Table 3-5). The Potomac and Patuxent rivers had the largest percentage of depauperate sites, failing for insufficient abundance or biomass. The Virginia mainstem also had a large percentage of depauperate sites, but this percentage was based on a comparatively small number of sites failing the restoration goals. The York and James rivers had the lowest percentages of depauperate sites. Low abundance, low biomass, and the level of widespread failure in most metrics necessary to classify a site as severely degraded would be expected on exposure to catastrophic events such as prolonged oxygen stress.

The upper Bay mainstem, Maryland eastern tributaries, James River, and York River, had excess abundance, excess biomass, or both in more than 27% of the failing sites (Table 3-6). Excess abundance and excess biomass are phenomena usually associated with eutrophic conditions and organic enrichment of the sediment in the absence of low dissolved oxygen stress.

Table 3-1. Summer trends in benthic community condition, 1985-2003. Trends were identified using the van Belle and Hughes (1984) procedure. Current mean B-IBI and condition are based on 2001-2003 values. Initial mean B-IBI and condition are based on 1985-1987 values. NS: not significant; (a): 1989-1991 initial condition; (b): 1995-1997 initial condition. Shaded areas highlight changes in trend or condition over those reported in 2002.

Station	Trend Significance	Median Slope (B-IBI units/yr)	Current Condition (2000-2002)	Initial Condition (1985-1987 unless otherwise noted)
Potomac River				
36	NS	0.00	2.17 Degraded	3.14 (Meets Goal)
40	NS	0.00	2.79 (Marginal)	2.80 (Marginal)
43	NS	0.00	3.67 (Meets Goal)	3.76 (Meets Goal)
44	p < 0.05	-0.04	1.84 (Severely Degraded)	2.80 (Marginal)
47	NS	0.00	3.53 (Meets Goal)	3.89 (Meets Goal)
51	p < 0.001	0.04	2.96 (Marginal)	2.43 (Degraded)
52	NS	0.00	1.22 (Severely Degraded)	1.37 (Severely Degraded)
Patuxent River				
71	NS	0.00	2.33 (Degraded)	2.59 (Degraded)
74	NS	0.00	3.44 (Meets Goal)	3.78 (Meets Goal)
77	p < 0.1	-0.07	3.40 (Meets Goal)	3.76 (Meets Goal)
79	NS	0.00	2.39 (Degraded)	2.75 (Marginal)
Choptank River				
64	NS	0.00	3.04 (Meets Goal)	2.78 (Marginal)
66	NS	0.00	2.73 (Marginal)	2.60 (Degraded)
Maryland Mainstem				
26	p < 0.01	0.03	3.93 (Meets Goal)	3.16 (Meets Goal)
24	NS	0.00	2.70 (Marginal)	3.04 (Meets Goal)
15	p < 0.05	0.03	2.70 (Marginal)	2.22 (Degraded)
06	p < 0.05	0.03	3.40 (Meets Goal)	2.56 (Degraded)
01	p < 0.05	0.03	3.59 (Meets Goal)	2.93 (Marginal)
Maryland Western Shore Tributaries				
22	NS	0.00	1.76 (Severely Degraded)	2.08 (Degraded)
23	NS	0.00	3.00 (Meets Goal)	2.49 (Degraded)
201	NS	0.00	1.49 (Severely Degraded)	1.10 (Severely Degraded) (a)
202	NS	0.00	1.89 (Severely Degraded)	1.40 (Severely Degraded) (a)
203	NS	0.02	2.07 (Degraded)	2.08 (Degraded) (b)
204	p < 0.05	-0.17	2.59 (Degraded)	3.67 (Meets Goal) (b)
Maryland Eastern Shore Tributaries				
29	p < 0.05	0.02	2.68 (Marginal)	2.38 (Degraded)
62	p < 0.05	-0.03	3.00 (Meets Goal)	3.42 (Meets Goal)
68	NS	0.00	3.40 (Meets Goal)	3.51 (Meets Goal)

Table 3-2. Summer trends in benthic community attributes at mesohaline stations 1985-2003. Monotonic trends were identified using the van Belle and Hughes (1984) procedure. ↑: Increasing trend; ↓: Decreasing trend. *: $p < 0.1$; **: $p < 0.05$; ***: $p < 0.01$; shaded trend cells indicate increasing degradation; unshaded trend cells indicate improving conditions; (a): trends based on 1989-2003 data; (b): trends based on 1995-2003 data; (c): attribute trend based on 1990-2003 data; (d): attributes are used in B-IBI calculations when species specific biomass is unavailable; NA: attribute is not part of the reported B-IBI. Blanks indicate no trend (not significant). See Appendix A for further detail.

Station	B-IBI	Abundance	Biomass	Shannon Diversity	Indicative Abundance	Sensitive Abundance	Indicative Biomass (c)	Sensitive Biomass (c)	Abundance Carnivore/Omnivores
Potomac River									
43					↑ ***	↓ ** (d)	NA		NA
44	↓ **	↓ ***	↓ ***			(d)	NA	↓ ***	NA
47					↑ **	↓ *** (d)	NA		NA
51	↑ ***		↓ ***	↑ ***	↓ ***	↑ ***	NA	NA	↑ **
52					(d)	(d)	↓ **		
Patuxent River									
71		↓ ***	↓ ***		↓ *** (d)	(d)			↑ ***
74		↑ ***	↓ **	↓ **	↑ *	↓ *** (d)	NA		NA
77	↓ *				↑ **	(d)	NA	↑ ***	NA
Choptank River									
64					(d)	↑ * (d)	↑ ***	↓ *	
Maryland Mainstem									
01	↑ **				↓ **	↑ ***	NA	NA	
06	↑ **	↑ **				↑ **	NA	NA	↑ *
15	↑ **	↑ **			↓ **		NA	NA	
24		↓ ***	↓ **	↓ ***	↓ ** (d)	↑ ** (d)			↑ ***
26	↑ ***					(d)	NA	↓ *	NA
Maryland Western Shore Tributaries									
22			↓ **	↓ *	↑ ***	(d)	NA		NA
23		↓ ***				↑ *** (d)	NA		NA
201(a)						(d)	NA		NA
202(a)			↑ **		↓ *	(d)	NA	↑ **	NA
204(b)	↓ **		↓ ***		↑ ** (d)	(d)	↑ ***	↓ **	
Maryland Eastern Shore Tributaries									
62	↓ **		↓ **	↓ ***	↓ *	(d)	NA		NA
68			↑ ***			↑ *** (d)	NA		NA

Table 3-3. Summer trends in benthic community attributes at oligohaline and tidal freshwater stations 1985-2003. Monotonic trends were identified using the van Belle and Hughes (1984) procedure. ↑: Increasing trend; ↓: Decreasing trend. *: $p < 0.1$; **: $p < 0.05$; ***: $p < 0.01$; shaded trend cells indicate increasing degradation; unshaded trend cells indicate improving conditions; (a): trends based on 1995-2003 data; NA: attribute not calculated. Blanks indicate no trend (not significant). See Appendix A for further detail.

Station	B-IBI	Abundance	Tolerance Score	Freshwater Indicative Abundance	Oligohaline Indicative Abundance	Oligohaline Sensitive Abundance	Tanypodinae to Chironomidae Ratio	Abundance Deep Deposit Feeders	Abundance Carnivore/Omnivores
Potomac River									
36					NA	NA	NA		NA
40				NA				NA	↑*
Patuxent River									
79		↑***		↓*	NA	NA	NA		NA
Choptank River									
66		↑***	↑**	NA			↑***	NA	↑**
Maryland Western Shore Tributaries									
203(a)				NA				NA	
Maryland Eastern Shore Tributaries									
29	↑**		↓***	NA	↓***			NA	↑**

Table 3-4. Estimated tidal area (km²) failing to meet the Chesapeake Bay benthic community restoration goals in the Chesapeake Bay, Maryland, Virginia, and each of the 10 sampling strata. In this table, the area of the mainstem deep trough is included in the estimates for the Severely Degraded portion of Chesapeake Bay, Maryland tidal waters, and Maryland mid-bay mainstem.

Region	Year	Severely Degraded	Degraded	Marginal	Total Failing	% Failing
Chesapeake Bay	1996	2,998	1,154	1,098	5,250	45.2
	1997	2,884	1,757	1,199	5,841	50.3
	1998	3,709	1,810	1,224	6,743	58.1
	1999	3,121	1,648	681	5,450	47.0
	2000	2,684	1,503	1,439	5,626	48.5
	2001	3,123	1,187	1,240	5,551	47.8
	2002	3,424	1,584	1,170	6,178	53.2
	2003	3,351	2,537	964	6,852	59.0
Maryland Tidal Waters	1994	2,684	1,152	497	4,332	66.5
	1995	2,872	605	182	3,659	58.6
	1996	2,614	700	155	3,469	55.6
	1997	2,349	697	483	3,529	56.5
	1998	2,663	1,016	623	4,302	68.9
	1999	2,423	1,137	374	3,935	63.0
	2000	2,455	1,137	236	3,828	61.3
	2001	2,313	582	644	3,538	56.7
	2002	2,444	713	928	4,086	65.4
	2003	2,571	1,288	228	4,086	65.4
Virginia Tidal Waters	1996	384	454	943	1,781	33.2
	1997	535	1,060	716	2,312	43.1
	1998	1,045	794	601	2,441	45.5
	1999	698	510	306	1,515	28.3
	2000	229	366	1,203	1,798	33.5
	2001	810	606	596	2,012	37.5
	2002	980	871	242	2,092	39.0
	2003	780	1,249	736	2,766	51.6
Potomac River	1994	793	330	0	1,123	60.7
	1995	510	153	51	714	56.0
	1996	714	51	0	765	60.0
	1997	561	204	102	867	68.0
	1998	561	510	102	1,173	92.0
	1999	663	153	102	918	72.0
	2000	612	255	0	867	68.0
	2001	612	357	51	1,020	80.0
	2002	561	204	153	918	72.0
	2003	867	153	0	1,020	80.0

Table 3-4. (Continued)						
Region	Year	Severely Degraded	Degraded	Marginal	Total Failing	% Failing
Patuxent River	1995	51	10	5	67	52.0
	1996	41	20	0	61	48.0
	1997	20	5	10	36	28.0
	1998	31	26	5	61	48.0
	1999	20	10	10	41	32.0
	2000	51	26	10	87	68.0
	2001	56	15	20	92	72.0
	2002	36	26	20	82	64.0
	2003	51	46	0	97	76.0
Maryland Upper Western Tributaries	1995	58	47	23	129	44.0
	1996	117	47	0	164	56.0
	1997	105	23	12	140	48.0
	1998	94	23	12	129	44.0
	1999	117	47	12	175	60.0
	2000	140	70	0	211	72.0
	2001	70	12	47	129	44.0
	2002	94	47	47	187	64.0
	2003	47	105	23	175	60.0
Maryland Eastern Tributaries	1995	107	128	0	235	44.0
	1996	21	150	21	192	36.0
	1997	43	64	21	128	24.0
	1998	21	64	64	150	28.0
	1999	43	150	86	278	52.0
	2000	64	150	21	235	44.0
	2001	128	64	86	278	52.0
	2002	64	107	64	235	44.0
	2003	128	214	0	342	64.0
Maryland Upper Bay Mainstem	1995	345	63	0	408	52.0
	1996	126	126	31	283	36.0
	1997	126	94	31	251	32.0
	1998	157	188	31	377	48.0
	1999	188	63	63	314	40.0
	2000	94	126	0	220	28.0
	2001	157	31	31	220	28.0
	2002	94	126	31	251	32.0
	2003	188	157	0	345	44.0

Region	Year	Severely Degraded	Degraded	Marginal	Total Failing	% Failing
Maryland Mid Bay Mainstem	1995	1,799	204	102	2,106	65.2
	1996	1,595	306	102	2,004	62.1
	1997	1,493	306	306	2,106	65.2
	1998	1,799	204	408	2,412	74.7
	1999	1,391	715	102	2,208	68.4
	2000	1,493	510	204	2,208	68.4
	2001	1,289	102	408	1,799	55.7
	2002	1,595	204	613	2,412	74.7
	2003	1,289	613	204	2,106	65.2
Virginia Mainstem	1996	165	330	824	1,318	32.0
	1997	165	824	659	1,648	40.0
	1998	824	330	494	1,648	40.0
	1999	494	165	165	824	20.0
	2000	0	165	1,154	1,318	32.0
	2001	494	330	494	1,318	32.0
	2002	659	659	165	1,483	36.0
	2003	494	824	659	1,977	48.0
Rappahannock River	1996	119	60	0	179	48.0
	1997	134	74	15	223	60.0
	1998	60	119	45	223	60.0
	1999	74	104	45	223	60.0
	2000	164	89	15	268	72.0
	2001	30	60	45	134	36.0
	2002	134	45	0	179	48.0
	2003	89	104	0	194	52.0
York River	1996	45	37	37	120	64.0
	1997	45	52	15	112	60.0
	1998	52	45	7	105	56.0
	1999	75	22	15	112	60.0
	2000	37	30	7	75	40.0
	2001	67	52	30	150	80.0
	2002	22	30	22	75	40.0
	2003	60	75	22	157	84.0
James River	1996	55	27	82	164	24.0
	1997	191	109	27	328	48.0
	1998	109	301	55	465	68.0
	1999	55	219	82	355	52.0
	2000	27	82	27	137	20.0
	2001	219	164	27	410	60.0
	2002	164	137	55	355	52.0
	2003	137	246	55	437	64.0

Table 3-5. Sites severely degraded (B-IBI < 2) and failing the restoration goals (scored at 1.0) for insufficient abundance, insufficient biomass, or both as a percentage of site failing the goals (B-IBI < 3), 1996 to 2003. Strata are listed in decreasing percent order of sites with insufficient abundance/biomass.

Stratum	Sites Severely Degraded		Sites Failing the Goals Due to Insufficient Abundance, Biomass, or Both	
	Number of Sites	As Percentage of Sites Failing the Goals	Number of Sites	As Percentage of Sites Failing the Goals
Potomac River	101	68.2	116	78.4
Patuxent River	60	55.0	85	78.0
Mid Bay Mainstem	64	55.2	84	72.4
West Tributaries	67	59.8	75	67.0
Virginia Mainstem	20	28.6	46	65.7
Rappahannock River	54	49.5	67	61.5
Upper Bay Mainstem	36	50.0	39	54.2
Eastern Tributaries	24	27.9	39	45.3
York River	54	41.2	46	35.1
James River	35	36.1	31	32.0

Table 3-6. Sites failing the restoration goals (scored at 1.0) for excess abundance, excess biomass, or both as a percentage of sites failing the goals (B-IBI < 3), 1996 to 2003. Strata are listed in decreasing percentage order.

Stratum	Number of Sites	As Percentage of Sites Failing the Goals
Eastern Tributaries	26	30.2
James River	28	28.9
York River	37	28.2
Upper Bay Mainstem	20	27.8
West Tributaries	22	19.6
Rappahannock River	20	18.3
Mid Bay Mainstem	17	14.7
Potomac River	21	14.2
Patuxent River	13	11.9
Virginia Mainstem	7	10.0




Figure 3-1. Results of probability-based benthic sampling of the Maryland Chesapeake Bay and its tidal tributaries in 2003. Each sample was evaluated in context of the Chesapeake Bay benthic community restoration goals.




Figure 3-2. Results of probability-based benthic sampling of the Virginia Chesapeake Bay and its tidal tributaries in 2003. Each sample was evaluated in context of the Chesapeake Bay benthic community restoration goals.

**Maryland Chesapeake Bay
Area Failing Restoration Goal**




Figure 3-3. Proportion of the Maryland Bay failing the Chesapeake Bay benthic community restoration goals from 1994 to 2003. The error bars indicate ± 1 standard error. The mainstem deep trough was sampled in 1994 and found to be mostly azoic; it is included in the severely degraded condition in 1994, but was excluded from sampling in subsequent years.

Chesapeake Bay 2003
Area Failing Restoration Goal




Figure 3-4. Proportion of the Chesapeake Bay, Maryland, Virginia, and the 10 sampling strata failing the Chesapeake Bay benthic community restoration goals in 2003. The error bars indicate ± 1 standard error.

Chesapeake Bay: Maryland
Stratum Area Failing Restoration Goal

3-16




Figure 3-5. Proportion of the Maryland sampling strata failing the Chesapeake Bay benthic community restoration goals, 1995 to 2003. The error bars indicate ± 1 standard error.

Chesapeake Bay: Virginia
Stratum Area Failing Restoration Goal

3-17




Figure 3-6. Proportion of the Virginia sampling strata failing the Chesapeake Bay benthic community restoration goals, 1996 to 2003. The error bars indicate ± 1 standard error.

**Chesapeake Bay
Area Failing Restoration Goal**




Figure 3-7. Proportion of the Chesapeake Bay failing the Chesapeake Bay benthic community restoration goals, 1996 to 2003. The error bars indicate ± 1 standard error.

4.0 DISCUSSION

Estimates of benthic community degradation for the Maryland Bay in 2003 were as high as those reported for 2002 (Llansó et al. 2003). However, the percent 'degraded' category was higher in 2003 than previously reported, and sites classified as 'marginally degraded' were fewer. Overall, 65% percent of the Maryland tidal waters failed the Chesapeake Bay benthic community restoration goals in 2003. Estimates of benthic community degradation for the Chesapeake Bay increased from 53% in 2002 to 59% in 2003, the largest estimate of degraded area since baywide monitoring began in 1996. The higher estimates for 2003 were associated with high flow conditions in the Bay, which were responsible for high nutrient and sediment run off, strong water column density stratification events, and widespread hypoxia. Wet conditions in 2003 are in contrast with below normal rainfall and low river flows in 2002 (Figure 4-1). The inter-annual changes in benthic condition we observe appear to be associated with these changes in hydrology (dry vs. wet years) and year-to-year fluctuations in dissolved oxygen concentrations. However, benthic community degradation in Chesapeake Bay continues to be large in any given year, and even though 2002 was a drought year, percent degradation for some monitoring strata was large (Llansó et al. 2003). We suggest that bay sediments are nutrient saturated. Excess organic matter retained in the sediments represents a rich source of food for benthic organisms, and primarily enhances growth and reproduction of the smaller, pollution tolerant forms. It will probably take sustained management efforts over an extended period of time to bring back a more balance community of benthic organisms and see significant bay-wide improvements in benthic condition.




Figure 4-1. Annual mean flow into Chesapeake Bay, 1937-2003. Unshaded area shows normal range of annual mean flows (25th to 75th percentile). Chart produced by the US Geological Survey and obtained from the USGS website at [http://md .water.usgs.gov/monthly/bay.html](http://md.water.usgs.gov/monthly/bay.html).

Fifty-one percent of the degraded Chesapeake Bay bottom in 2003 (3,501 km²) was marginally to moderately impaired. In the Maryland portion of the Bay, 37% of the degraded bottom (1,516 km²) was marginally to moderately impaired. Of the additional 2,571 km² of Maryland Bay bottom supporting severely degraded benthic communities, 676 km² were located in the deep (>12m) mainstem that is perennially anoxic and probably beyond the scope of present mitigation efforts. No obvious trends in the percentage of area with marginal or moderate degradation were observed over the time series.

The estimates of degraded area for regions measured in multiple years were generally similar between years, with most estimates included within the confidence interval of other years. Some exceptions can be explained by the clumping of the random sites in either deep areas that are perennially hypoxic (e.g., the exceptionally high estimate of degraded area for the Potomac River in 1998) or shallow areas that are not typically affected by summer hypoxia (e.g., the low estimate of degraded area for the Patuxent River in 1997 and 1999). In addition, inter-annual variability in flow patterns influences water quality and benthic community condition. High spring flows, for example, have been theorized to cause earlier and spatially more extensive stratification within the Bay, leading to more extensive hypoxia (Tuttle et al. 1987). Patterns of degradation between years, although subtle, were in the direction expected from abnormally strong spring freshets.

Below we discuss the patterns of degradation and sources of stress affecting benthic communities in each of the Maryland Bay six strata (see Figure 2-4) as inferred from the results of the water quality and the benthic monitoring programs.

4.1 PATUXENT RIVER

Benthic degradation in the Patuxent River is mainly related to adverse effects from low dissolved oxygen (DO). The intensity of summer hypoxic events varies annually, and this variability is reflected in the B-IBI. As indicated in Figure 4-2, there is a positive relationship between the percentage of samples failing the restoration goals (B-IBI scores less than 3.0) and summer hypoxia, expressed as percent observations in the mesohaline Patuxent River with bottom DO concentrations below 2 mg/L, as measured at water quality monitoring stations, June through September. Hypoxia in 2003 was severe, and thus a majority of samples in the Patuxent River failed the restoration goals. A strong relationship was also observed for the 1995-2002 time series when the average DO concentration measured at the time of the benthic sampling was plotted against the percentage of samples failing the restoration goals (Llansó et al. 2003). That relationship explained 76% of the variability. With the addition of the 2003 data, the strength of the relationship decreased substantially (Figure 4-3). This was due to higher DO concentrations in the lower Patuxent River at the time of the benthic sampling (late August 2003) than during the preceding month. Bay-wide, hypoxia was more extensive and severe in July, and large mortality of benthic organisms probably occurred at that time, with little recovery of the community in the following weeks. Also, many of the 2003 benthic sites in the lower Patuxent River were located in shallow water above the pycnocline, and these locations exhibited relatively high DO concentrations in late August 2003.




Figure 4-2. Relationship of benthic index of biotic integrity to percent dissolved oxygen observations below 2 mg/L (June-September) in the mesohaline Patuxent River. Each point represents a different year, 1995-2003. Dissolved oxygen data are fortnight near-bottom observations from Chesapeake Bay Water Quality Monitoring Program stations RET1.1, and LE1.1 through LE1.4.




Figure 4-3. Relationship of benthic index of biotic integrity to dissolved oxygen concentration at the time of benthic sample collection in the mesohaline Patuxent River. Each point represents a different year, 1995-2003.

One factor linked to hypoxia is the amount of decaying organic matter from phytoplankton blooms. The lower Patuxent River suffers from poor water clarity and high algal concentrations. Years with large phytoplankton blooms are likely to result in more extensive hypoxia and increased benthic degradation. A positive association between the percentage of samples with severely degraded benthic condition and average chlorophyll *a* concentrations in the lower Patuxent River was still evident with the addition of the 2003 data (Figure 4-4). There were strong relationships for average chlorophyll concentrations below the pycnocline for quarter 2 (April-June), quarter 3 (July-September), and the combined quarters 1-3. Above the pycnocline, chlorophyll concentrations in the lower Patuxent River were highest in 2003, with maximum observed concentrations greater than 700 µg/L.




Figure 4-4. Relationship of benthic index of biotic integrity to average chlorophyll *a* concentration in the mesohaline Patuxent River. Each point represents a different year, 1995-2003. Chlorophyll data are below pycnocline, April through June fortnight observations from Chesapeake Bay Water Quality Monitoring Program stations RET1.1, and LE1.1 through LE1.4.

Benthic community status at three of the four fixed monitoring stations in the Patuxent River showed declines in condition in 2003. Station 77 (Holland’s Cliff) showed improvement. The most severe declines occurred at Station 71 (Broomes Island) and Station 79 (tidal freshwater at Lyons Creek). The low B-IBI scores at the fixed monitoring stations were most likely associated with the severe hypoxia and high river flow conditions of 2003. Long-term trends at these stations, however, continued to give signals of

recovery. At Station 77, the magnitude of the degrading trend (declining B-IBI) continued to diminish. In the last three years, recovery at this station was associated with increases in biomass. The biomass metric is now scoring in the good range, reflecting increasing densities of the bivalves *Macoma balthica* and *Rangia cuneata*, and the crustacean *Cyathura polita*. This trend, however, should be viewed with caution, as flow-induced changes in salinity typically limit the distribution of bivalves in the Chesapeake Bay (Holland et al. 1987) and are likely to play a major role in structuring benthic communities in transitional salinity regions.

In addition to the positive changes in benthic community condition noted for Station 77, a significant degrading trend through 2001 at Station 71 (Broomes Island) disappeared with the addition of the 2002 data. Poor benthic condition at this station in 2003 did not change that result. However, variable annual low DO events are likely to influence direction in B-IBI trends at Station 71. We note continuing degrading trends in total community abundance and biomass (see Table 3-2).

4.2 POTOMAC RIVER

The Potomac River has one of the highest areas with degraded benthic community in the Chesapeake Bay. Much of the problem in the Potomac River is severe oxygen depletion in the lower deep mainstem. Over the period 1996-2003, this stratum had the highest percentage of sites failing the restoration goals because of insufficient abundance or biomass. Unlike with the Patuxent River, no significant relationship was observed when the percent samples failing the restoration goals was plotted against the percentage of observations with DO concentrations below 2 mg/L, June through September. This is because hypoxia in the Potomac River is a perennial problem that affects waters below the pycnocline, with little inter-annual variability. In 2003, 64% of the bottom DO readings at the Potomac mesohaline water quality monitoring stations were below 2 mg/L. Although this percentage was not much higher than those reported for previous years, 59% of the observations were below 1 mg/L, the highest value since 1994.

A relationship was observed when examining the average DO concentration of benthic sites in relation to percent samples failing the restoration goals (Figure 4-5). Hypoxic events tend to be long lasting in the main stem of the Potomac River and thus, August DO concentrations continued to be low during this month. All (100%) of the sites sampled in the mesohaline Potomac River in 2003 failed the restoration goals. Relationships between the B-IBI and DO in the Potomac River, however, are best explored as a function of depth. The frequency of low DO events in the Potomac River is strongly associated with water depth (Figure 4-6), and so is the probability of observing severely degraded benthos (Figure 4-7). Unlike with the Patuxent River, relationships between the B-IBI and chlorophyll concentrations in the Potomac River are not particularly strong.




Figure 4-5. Relationship of benthic index of biotic integrity to dissolved oxygen concentration at the time of benthic sample collection in the mesohaline Potomac River. Each point represents a different year, 1994-2003.




Figure 4-6. Relationship between percent DO observations below 2 mg/L and water depth in the mesohaline Potomac River




Figure 4-7. Probability of observing severely degraded benthos (B-IBI ≤ 2.0) as a function of water depth in the mesohaline Potomac River. A logistic regression model was used to obtain the probabilities.

Of the seven fixed monitoring stations in the Potomac River, only two showed trends in the B-IBI. Station 44 at Morgantown exhibited a degrading trend (significantly declining B-IBI), and Station 51 in shallow water near St. Clements Island exhibited an improving trend (significantly increasing B-IBI). The trend at Station 44 was new in 2002 and has become more pronounced with the addition of the 2003 data. Degrading trends in abundance and biomass contributed to the observed B-IBI trend. Shannon diversity also declined. Station 44 is on the slope of the deep channel (11-17 m) of the Potomac River and may be affected by tilts of the pycnocline bringing episodic fluctuations in DO and salinity. These are likely to exert severe stress on the benthic community, especially if abrupt changes in flow or salinity occur during the benthic reproductive season. For example, in May 1998 (a wet year) salinity at Station 44 was 1.5 psu, while in May 2002 (a dry year), salinity was 21 psu. The long-term summer salinity average is in the low mesohaline range. DO at the time of the benthic sampling was relatively high in 1999, 2000, and 2002 (5-7 mg/L), just above 2 mg/L in 1996, 1997, and 2001, and below 2 mg/L in 2003.

The improving trend at Station 51 was reinforced with the 2003 data, and was due to significant increases in diversity, pollution-sensitive abundance, and carnivore-omnivore abundance, and to significant decreases in pollution-indicative abundance. These changes may indicate improving water quality conditions in the lower shallow Potomac River. Total community biomass, however, has been steadily decreasing since the 1980s.

4.3 UPPER WESTERN TRIBUTARIES

Benthic community impairment in the upper western tributaries of the Bay was high in 2003, with 60% of the area exhibiting degraded condition. The western tributaries suffer from various types of pollution, including toxic contamination, low dissolved oxygen, excess phytoplankton growth, lack of water clarity, and nutrient runoff, but these factors vary greatly among systems and the stress to the benthic communities varies accordingly. Patterns described in previous years were reinforced with the addition of the 2003 data. Results indicate good agreement between the status and trends for water quality parameters and the benthic community condition.

Benthic community condition is severely degraded in the upper part of the Patapsco River estuary, above the Francis Scott Key Bridge and at sites in Curtis Creek, Stony Creek, and along the deep channel south of Sparrows Point, areas that are affected by very low DO concentrations and by toxic contamination. Excess abundance, indicating eutrophic conditions, is common in the lower portion of the estuary in areas that are not affected by hypoxia. In contrast with the Patapsco River, summer DO status is good in the Back River (Tidal Monitoring and Analysis Workgroup). Back River mesohaline stations show excess phytoplankton growth and poor Secchi depth. The Back River shows moderately degraded benthic condition with total densities of organisms that are either within the good range or in excess of reference conditions, in agreement with pollution related to excess algal growth and high particulate organic deposition.

Good benthic community condition in the Middle River is consistent with observations of good water quality status for this river. The Gunpowder River shows patterns of benthic community degradation over a large area. In contrast, the Bush River shows degradation in the upper reaches of the estuary. Degraded sites in the Bush River were numerically dominated by pollution tolerant organisms, mostly tubificid oligochaetes. This is consistent with poor water quality for chlorophyll *a* and Secchi depth in this region of the river (Tidal Monitoring and Analysis Workgroup).

To the south, the Magothy River exhibited degradation in the upper two thirds of the estuary. Patterns of degradation appeared to respond to a mixture of over-enrichment and hypoxia. This is consistent with excess algal abundance and observations of low DO concentrations at shallow, upstream water quality monitoring stations. In the Severn River, benthic community degradation was restricted to the upper portion of the estuary, above the long-term water quality monitoring station (WT7.1). Sites in this region of the river had few organisms or were azoic, consistent with severe hypoxia or anoxia problems identified during the assessments conducted in 2002 using DATAFLOW technology (see <http://mddnr.chesapeakebay.net/eyesonthebay/index.cfm>). The fixed benthic monitoring station in the Severn River (Station 204) showed a degrading trend (significantly declining B-IBI) in 2003. This new trend may be signaling an increase in the extent of the low DO area. The South, Rhode, and West Rivers exhibited about 50% degradation predominately due to excess abundance indicative of over-enrichment.

4.4 EASTERN TRIBUTARIES

The Maryland eastern tributaries usually have some of the smallest extent of degraded area in the Chesapeake Bay. However, degradation was exceptionally high in 2003 with 64% of the area (342 km²) failing the restoration goals. Degradation affected predominately the lower Chester River and the rivers emptying in Tangier Sound, with the exception of the Wicomico River. A majority of the sites with failing B-IBI in the Chester River were concentrated in the lower portion of the river, around Eastern Neck Island. Poor benthic community condition in this region could not be attributed to stress from low DO. Fifty percent of the sites in this region exhibited excess abundance of organisms, which is consistent with degrading trends in chlorophyll concentrations and water clarity. A fixed station (Station 68) located mid-river above the region where a majority of the random samples fail the B-IBI, had good status (meets goal) and exhibited no trend.

Maryland eastern tributaries have high agricultural land use, high nutrient input, high chlorophyll values but low frequencies of low dissolved oxygen events (Dauer et al. 2000). A high incidence of failure of restoration goals due to excess abundance of organisms is observed for these tributaries. However, in the lower eastern shore basin, low biomass relative to reference conditions is a problem, particularly in the Manokin River and Tangier Sound. Overall, nearly half of the sites in the lower eastern tributaries (Nanticoke, Wicomico, Manokin, Big Annemessex, and Pocomoke) had low biomass. The major problem affecting water quality in the lower eastern shore basin is high sediment loads, which may reduce the amount of food that is available from the water column to the benthos. High river flows in 2003 may have exacerbated the high sediment load problem.

The fixed long-term monitoring station in the Nanticoke River (Station 62) exhibited a degrading trend in the B-IBI. However, the strength of the B-IBI trend and of declining trends in biomass and Shannon diversity, diminished with the addition of the 2003 data. Benthic community status at Station 62 now meets the goal. High sediment and nutrient loads are major problems in the Nanticoke River. The fixed station in the Elk River (Station 29) exhibited a significant, positive trend in the B-IBI, but marginally degraded benthic community status in 2003. The improving trend was associated with a decrease in the abundance of tubificid oligochaetes and an overall increase in the densities of the bivalve *Rangia cuneata*. Although improving trends in this region are reported for nutrients, chlorophyll *a*, and sediment concentrations, benthic community dynamics in the Elk River are likely to be influenced by patterns in river flow and the associated salinity fluctuations. Thus, wet years may enhance the abundance of tubificid oligochaetes.

4.5 MARYLAND MID BAY AND UPPER BAY MAINSTEMS

Low DO events are common and severe in the mid-bay Maryland mainstem (Dauer et al. 2000). Anoxia is a common feature of the mid-bay deep channel. The Maryland mainstem stratum has the largest extent of severely degraded benthic community condition in the Bay. Three fixed stations are located in shallow, sandy habitats of the mainstem (Stations 01, 06, and 15). These stations showed significant improving trends

in the B-IBI in 2003, and marginal (Station 15) or good (Stations 01 and 06) benthic community status.

The upper Maryland mainstem receives discharges from the Susquehanna River; therefore, water quality in this region is a good indicator of inputs from the Susquehanna River watershed. A high incidence of failure of restoration goals due to excess abundance or biomass of organisms is a common feature in the upper bay. This is indicative of effects on benthos resulting from nutrient enrichment. However, Station 26 further to the south showed a significant positive trend in the B-IBI and good benthic community status, which did not change with the more extensive hypoxia affecting the bay in 2003.

4.6 VIRGINIA TRIBUTARIES

All of the Virginia tributaries experienced increases in the extent of benthic community degradation in 2003. Mainstem hypoxia in the bay covered extensive areas of the lower bay in July 2003 (Tidal Monitoring and Analysis Workgroup). This is likely to have been associated with the higher than usual benthic community degradation in the Virginia mainstem. The York River exhibited the largest percent of area degraded. The York rivers does not normally experience hypoxia, except for periods of intermittent hypoxia associated with spring-neap tidal cycles (Haas 1977). Sites in the deep channel of the lower York River had low DO concentrations in 2003, but total community abundance was not particularly low. Many sites throughout the York River exhibited excess abundance of organisms, a condition more often associated with nutrient enrichment. Physical disturbance of the sediments associated with strong erosional and depositional events is also known to structure benthic communities in the York River (Schaffner et al. 2002). These events were documented through radioisotope dating of sediments and were associated with tidal exchange and river flow.

In the James River, patterns in benthic community condition among years can be partially explained by the clumping of sites in areas with local contamination problems. Because pollution sources are spatially variable in the James River stratum, comparisons in patterns of benthic community condition should be interpreted with caution and include assessments at various spatial scales of variability (Dauer and Llansó 2003). Patterns of degradation in the James River are driven by serious sediment contamination problems concentrated in the Elizabeth River (Dauer and Llansó 2003).

4.7 CONCLUSIONS

Baywide estimates of degradation in 2003 were high and in tune with high river flows and extensive hypoxia. However, positive trends in benthic community condition continued to be detected at some fixed long-term monitoring stations, most notably in the Patuxent River estuary. Local areas with identifiable point sources of pollution may be the first ones to respond to pollution abatement and are more likely to show recovery at fixed stations.

The probability-based estimates developed for this report are the result of reviews conducted jointly by the Maryland and Virginia Chesapeake Bay benthic monitoring programs. A program review in 1996 examined program objectives, analysis techniques, and power to detect trends. One objective that emerged from the program review process was a goal of producing a baywide area estimate of degraded benthic communities with known and acceptable uncertainty. That goal is now an inherent part of benthic monitoring activities in Chesapeake Bay.

Baywide estimates are dependent on fully validated thresholds for assessing the condition of the benthic community in each sample collected. The thresholds were established and validated by Ranasinghe et al. (1994a) and updated by Weisberg et al. (1997). The B-IBI and the stratified random sampling design allow a validated, unambiguous approach to characterizing conditions in the Chesapeake Bay. The Chesapeake Bay B-IBI has been shown by Alden et al. (2002) to be sensitive, stable, robust, and statistically sound. The B-IBI is also applicable to a wide range of habitats, from tidal freshwater mud to polyhaline sand in the Chesapeake Bay, and this is an important and useful feature of the index because it allows characterization of local gradients of pollution and conditions across habitats. A study to develop diagnostic tools that differentiate between low dissolved oxygen impacts on benthos and those from toxic contamination was recently conducted by Dauer et al. (2002) and further augmented the usefulness of the B-IBI to management.

Although a continuing evolution of the B-IBI may lead to changes in estimates of the area of the Bay meeting the restoration goals, these revisions should amount to fine-tuning and not to significant changes in the estimates. One strength of the probability-based sampling element is that the amount of area meeting the goals can be recalculated as the index continues to be improved, so that trends in the area meeting the goals can be compared in a consistent and rigorous fashion.

5.0 REFERENCES

- Alden, R.W. III, D.M. Dauer, J.A. Ranasinghe, L.C. Scott, and R.J. Llansó. 2002. Statistical verification of the Chesapeake Bay benthic index of biotic integrity. *Environmetrics* 13:473-498.
- Alden, R. W. III, J.A. Ranasinghe, L.C. Scott, R.J. Llansó, and D. M. Dauer. 2000. B-IBI Phase 3: Optimization of the benthic index of biotic integrity. Prepared for the Maryland Department of Natural Resources by Versar, Inc., Columbia, MD.
- Alden, R.W. III, S.B. Weisberg, J.A. Ranasinghe, and D.M. Dauer. 1997. Optimizing temporal sampling strategies for benthic environmental monitoring programs. *Marine Pollution Bulletin* 34:913-922.
- Baird, D. and R.E. Ulanowicz. 1989. The seasonal dynamics of the Chesapeake Bay Ecosystem. *Ecological Monographs* 59:329-364.
- Boicourt, W.C. 1992. Influences of circulation processes on dissolved oxygen in the Chesapeake Bay. Pages 7-59. *In*: D.E. Smith, M. Leffler, and G. Mackiernan (eds.), *Oxygen Dynamics in the Chesapeake Bay: A Synthesis of Recent Results*. Maryland Sea Grant Program, College Park, MD.
- Boynton, W.R. and W.M. Kemp. 2000. Influence of river flow and nutrient loads on selected ecosystem processes: A synthesis of Chesapeake Bay data. Pages 269-298. *In*: J. E. Hobbie, ed., *Estuarine Science: A Synthetic Approach to Research and Practice*. Island Press, Washington, D.C.
- Dauer, D.M. 1993. Biological criteria, environmental health and estuarine macrobenthic community structure. *Mar. Pollution Bulletin* 26:249-257.
- Dauer, D.M. and W.G. Conner. 1980. Effects of moderate sewage input on benthic polychaete populations. *Estuarine, Coastal, and Marine Science* 10:335-346.
- Dauer, D.M., M.F. Lane, and R.J. Llansó. 2002. Development of diagnostic approaches to determine sources of anthropogenic stress affecting benthic community condition in the Chesapeake Bay. Prepared for U.S. Environmental Protection Agency, Chesapeake Bay Program Office, by Department of Biological Sciences, Old Dominion University, Norfolk, VA.
- Dauer, D.M. and R.J. Llansó. 2003. Spatial scales and probability based sampling in determining levels of benthic community degradation in the Chesapeake Bay. *Environmental Monitoring and Assessment* 81:175-186.
- Dauer, D.M., M.W. Luchenback, and A.J. Rodi, Jr. 1993. Abundance biomass comparisons (ABC method): Effects of an estuary gradient, anoxic/hypoxic events, and contaminated sediments. *Marine Biology* 116:507-518.

- Dauer, D.M., J.A. Ranasinghe, and S.B. Weisberg. 2000. Relationships between benthic community condition, water quality, sediment quality, nutrient loads, and land use patterns in Chesapeake Bay. *Estuaries* 23:80-96.
- Dauer, D.M., A.J. Rodi, Jr., and J.A. Ranasinghe. 1992. Effects of low dissolved oxygen events on the macrobenthos of the lower Chesapeake Bay. *Estuaries* 15:384-391.
- Dennison, W.C., R.J. Orth, K.A. Moore, J.C. Stevenson, V. Carter, S. Kollar, P.W. Bergstrom, and R.A. Batiuk. 1993. Assessing water quality with submersed aquatic vegetation. Habitat requirements as barometers of Chesapeake Bay health. *BioScience* 43:86-94.
- Diaz, R.J. and R. Rosenberg. 1995. Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanography and Marine Biology Annual Review* 33:245-303.
- Diaz, R.J. and L.C. Schaffner. 1990. The functional role of estuarine benthos. Pages 25-56. In: M. Haire and E. C. Chrome, eds., *Perspectives on the Chesapeake Bay*, Chapter 2. Chesapeake Research Consortium, Gloucester Point, VA. CBP/TRS 41/90.
- Flemer, D.A., G.B. Mackiernan, W. Nehlsen, and V.K. Tippie. 1983. Chesapeake Bay: A profile of environmental change. U.S. Environmental Protection Agency, Washington, DC.
- Frithsen, J. 1989. The benthic communities within Narragansett Bay. An assessment for the Narragansett Bay Project by the Marine Ecosystems Research Laboratory, Graduate School of Oceanography, University of Rhode Island, Narragansett, RI.
- Gray, J.S. 1979. Pollution-induced changes in populations. *Philosophical Transactions of the Royal Society of London* B286:545-561.
- Haas, L.W. 1977. The effect of the spring-neap tidal cycle on the vertical salinity structure of the James, York and Rappahannock Rivers, Virginia, U.S.A. *Estuarine, Coastal, and Marine Science* 5:485-496.
- Holland, A.F., N.K. Mountford, M.H. Hiegel, K.R. Kaumeyer, and J.A. Mihursky. 1980. The influence of predation on infaunal abundance in upper Chesapeake Bay. *Marine Biology* 57:221-235.
- Holland, A.F., N.K. Mountford, and J.A. Mihursky. 1977. Temporal variation in the upper bay mesohaline benthic communities: 1. The 9-m mud habitat. *Chesapeake Science* 18:370-378.

- Holland, A.F., A.T. Shaughnessy, and M.H. Hiegel. 1987. Long-term variation in mesohaline Chesapeake Bay macrobenthos: Spatial and temporal patterns. *Estuaries* 3:227-245.
- Holland, A.F., A.T. Shaughnessy, L.C. Scott, V.A. Dickens, J.A. Ranasinghe, and J.K. Summers. 1988. Long-term benthic monitoring and assessment program for the Maryland portion of Chesapeake Bay (July 1986-October 1987). Prepared for Power Plant Research Program, Department of Natural Resources and Maryland Department of the Environment by Versar, Inc., Columbia, MD.
- Holland, A.F., A.T. Shaughnessy, L.C. Scott, V.A. Dickens, J. Gerritsen, and J.A. Ranasinghe. 1989. Long-term benthic monitoring and assessment program for the Maryland portion of Chesapeake Bay: Interpretive report. Prepared for the Maryland Dept. of Natural Resources by Versar, Inc., Columbia, MD. CBRM-LTB/EST-2.
- Homer, M. and W.R. Boynton. 1978. Stomach analysis of fish collected in the Calvert Cliffs region, Chesapeake Bay-1977. Final report prepared for the Maryland Power Plant Siting Program by the University of Maryland, Chesapeake Biological Laboratory, Solomons, MD. UMCEES 78-154-CBL.
- Homer, M., P.W. Jones, R. Bradford, J.M. Scolville, D. Morck, N. Kaumeyer, L. Hoddaway, and D. Elam. 1980. Demersal fish food habits studies near Chalk Point Power Plant, Patuxent estuary, Maryland, 1978-1979. Prepared for the Maryland Department of Natural Resources, Power Plant Siting Program, by the University of Maryland, Center for Environmental and Estuarine Studies, Chesapeake Biological Laboratory, Solomons, MD. UMCEES-80-32-CBL.
- Llansó, R.J. 1992. Effects of hypoxia on estuarine benthos: The lower Rappahannock River (Chesapeake Bay), a case study. *Estuarine, Coastal, and Shelf Science* 35:491-515.
- Llansó, R.J., D.M. Dauer, J.H. Vølstad, and L.C. Scott. 2003. Application of the benthic index of biotic integrity to environmental monitoring in Chesapeake Bay. *Environmental Monitoring and Assessment* 81:163-174.
- Llansó, R.J., L.C. Scott, and F.S. Kelley. 2003. Chesapeake Bay water quality monitoring program: Long-term benthic monitoring and assessment component, Level 1 Comprehensive Report (July 1984-December 2002). Prepared for the Maryland Department of Natural Resources by Versar, Inc., Columbia, MD.
- Malone, T.C. 1987. Seasonal oxygen depletion and phytoplankton production in Chesapeake Bay: Preliminary results of 1985-86 field studies. Pages 54-60. *In*: G.B. Mackiernan, ed., Dissolved Oxygen in the Chesapeake Bay: Processes and Effects. Maryland Sea Grant, College Park, MD.

- Malone, T.C., L.H. Crocker, S.E. Pile, and B.W. Wendler. 1988. Influences of river flow on the dynamics of phytoplankton production in a partially stratified estuary. *Marine Ecology Progress Series* 48:235-249.
- National Research Council (NRC). 1990. Managing Troubled Waters: The Role of Marine Environmental Monitoring. National Academy Press, Washington, DC.
- Officer, C.B., R.B. Biggs, J.L. Taft, L.E. Cronin, M.A. Tyler, and W.R. Boynton. 1984. Chesapeake Bay anoxia: Origin, development, and significance. *Science* 223:22-27.
- Pearson, T.H. and R. Rosenberg. 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanography and Marine Biology Annual Review* 16:229-311.
- Ranasinghe, J.A., L.C. Scott, and S.B. Weisberg. 1993. Chesapeake Bay water quality monitoring program: Long-term benthic monitoring and assessment component, Level 1 Comprehensive Report (July 1984-December 1992). Prepared for Maryland Department of the Environment and Maryland Department of Natural Resources by Versar, Inc., Columbia, MD.
- Ranasinghe, J.A., S.B. Weisberg, D.M. Dauer, L.C. Schaffner, R.J. Diaz, and J.B. Frithsen. 1994a. Chesapeake Bay Benthic Community Restoration Goals. Prepared for the U.S. Environmental Protection Agency Chesapeake Bay Program Office, the Governor's Council on Chesapeake Bay Research Fund, and the Maryland Department of Natural Resources by Versar, Inc., Columbia, MD.
- Ranasinghe, J.A., S.B. Weisberg, J. Gerritsen, and D.M. Dauer. 1994b. Assessment of Chesapeake Bay benthic macroinvertebrate resource condition in relation to water quality and watershed stressors. Prepared for The Governor's Council on Chesapeake Bay Research Fund and the Maryland Department of Natural Resources by Versar, Inc., Columbia, MD.
- Ritter, C. and P.A. Montagna. 1999. Seasonal hypoxia and models of benthic response in a Texas Bay. *Estuaries* 22:7-20.
- Schaffner, L.C., T.M. Dellapenna, E.K. Hinchey, C.T. Friedrichs, M.T. Neubauer, M.E. Smith, and S.A. Kuehl. 2002. Physical energy regimes, seabed dynamics and organism-sediment interactions along an estuarine gradient. Pages 159-180. *In: J.Y. Aller, S.A. Woodin, and R.C. Aller, eds., Organism-Sediment Interactions.* University of South Carolina Press, Columbia, SC.

- Scott, L.C., A.F. Holland, A.T. Shaughnessy, V. Dickens, and J.A. Ranasinghe. 1988. Long-term benthic monitoring and assessment program for the Maryland portion of Chesapeake Bay: Data summary and progress report. Prepared for Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, and Maryland Department of the Environment by Versar, Inc., Columbia, MD. PPRP-LTB/EST-88-2.
- Seliger, H.H., J.A. Boggs, and W.H. Biggley. 1985. Catastrophic anoxia in the Chesapeake Bay in 1984. *Science* 228:70-73.
- Sen, P.K. 1968. Estimates of the regression coefficient based on Kendall's tau. *Journal of the American Statistical Association* 63:1379-1389.
- Tuttle, J.H., R.B. Jonas, and T.C. Malone. 1987. Origin, development and significance of Chesapeake Bay anoxia. Pages 443-472. *In*: S.K. Majumdar, L.W. Hall, Jr., and H.M. Austin, eds., Contaminant Problems and Management of Living Chesapeake Bay Resources. Pennsylvania Academy of Science, Philadelphia, PA.
- van Belle, G. and J.P. Hughes. 1984. Nonparametric tests for trend in water quality. *Water Resources Research* 20:127-136.
- Versar, Inc. 1999. Versar Benthic Laboratory Standard Operating Procedures and Quality Control Procedures. Versar, Inc., Columbia, MD.
- Virnstein, R.W. 1977. The importance of predation of crabs and fishes on benthic infauna in Chesapeake Bay. *Ecology* 58:1199-1217.
- Warwick, R.M. 1986. A new method for detecting pollution effects on marine macrobenthic communities. *Marine Biology* 92:557-562.
- Weisberg, S.B., J.A. Ranasinghe, D.M. Dauer, L.C. Schaffner, R.J. Diaz, and J.B. Frithsen. 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. *Estuaries* 20:149-158.
- Wilson, J.G. and D.W. Jeffrey. 1994. Benthic biological pollution indices in estuaries. Pages 311-327. *In*: J.M. Kramer, ed., Biomonitoring of Coastal Waters and Estuaries. CRC Press, Boca Raton, FL.

APPENDIX A

**FIXED SITE COMMUNITY ATTRIBUTE
1985-2003 TREND ANALYSIS RESULTS**

Appendix Table A-1. Summer trends in benthic community attributes at mesohaline stations 1985-2003. Shown is the median slope of the trend. Monotonic trends were identified using the van Belle and Hughes (1984) procedure. Shaded cells indicate increasing degradation; unshaded cells indicate improving conditions; (a): trends based on 1989-2003 data; (b): trends based on 1995-2003 data; (c): attribute trend based on 1990-2003 data; (d): attributes are used in B-IBI calculations when species specific biomass is unavailable; (e): attribute and trend are not part of the reported B-IBI.

Station	B-IBI	Abundance	Biomass	Shannon Diversity	Indicative Abundance	Sensitive Abundance	Indicative Biomass (c)	Sensitive Biomass (c)	Abundance Carnivore/Omnivores
Potomac River									
43	0.00	-47.06	-0.72	0.003	0.34	-1.80 (d)	0.005 (e)	-0.07	-0.13 (e)
44	-0.04	-45.00	-0.13	0.02	-0.30	-0.24(d)	0.01 (e)	-3.62	0.74 (e)
47	0.00	-13.33	0.70	0.02	0.28	-1.15 (d)	-0.001 (e)	-0.83	-0.36 (e)
51	0.04	0.00	-0.19	0.02	-1.20	0.65	0.11 (e)	-0.58 (e)	0.65
52	0.00	-3.25	-0.00	0.00	0.00 (d)	0.00 (d)	0.00	0.00	0.00
Patuxent River									
71	0.00	-49.23	-0.08	0.01	-2.09 (d)	0.00 (d)	-1.20	0.00	1.11
74	0.00	199.19	-0.85	-0.03	0.25	-1.42 (d)	0.00 (e)	-0.05	-0.60 (e)
77	-0.04	41.79	-0.12	0.004	1.17	-0.38 (d)	-3.66 (e)	7.98	-0.38 (e)
Choptank River									
64	0.00	30.62	0.03	0.02	-0.23 (d)	0.67 (d)	0.18	-1.69	-0.14
Maryland Mainstem									
01	0.03	15.00	0.03	-0.002	-0.49	1.06	-0.11 (e)	0.63 (e)	0.61
06	0.03	42.86	0.006	0.02	-0.17	1.36	0.00 (e)	-0.04 (e)	0.94
15	0.03	44.00	-0.02	0.003	-0.96	0.13	-0.003 (e)	0.62 (e)	0.29
24	0.00	-55.31	-0.20	-0.04	-0.52 (d)	0.49 (d)	-0.003	0.03	1.32
26	0.03	20.00	-0.04	0.02	0.00	0.20 (d)	0.00 (e)	-0.04	0.50 (e)
Maryland Western Shore Tributaries									
22	0.00	-23.27	-0.04	-0.04	2.22	0.00 (d)	0.99 (e)	-0.03	-0.65 (e)
23	0.00	-90.91	-0.00	0.002	0.17	0.53 (d)	-0.014 (e)	1.11	0.38 (e)
201(a)	0.00	-12.50	-0.001	0.00	0.00	0.00 (d)	4.72 (e)	0.00	0.00 (e)
202(a)	0.00	15.15	0.002	0.06	-1.29	0.00 (d)	-0.15 (e)	0.00	1.25 (e)
204(b)	-0.17	-34.23	-0.46	0.005	5.01 (d)	-0.32 (d)	0.23	-4.56	-0.40
Maryland Eastern Shore Tributaries									
62	-0.03	69.33	-0.05	-0.05	-0.10	-0.29 (d)	0.00 (e)	-0.19	-0.28 (e)
68	0.00	62.32	0.87	0.01	0.05	1.65 (d)	-0.002 (e)	0.001	0.98 (e)

Appendix Table A-2. Summer trends in benthic community attributes at oligohaline and tidal freshwater stations 1985-2003. Shown is the median slope of the trend. Monotonic trends were identified using the van Belle and Hughes (1984) procedure. Shaded cells indicate increasing degradation; unshaded cells indicate improving conditions; (a): trends based on 1989-2003 data; NA: attribute not calculated.

Station	B-IBI	Abundance	Tolerance Score	Freshwater Indicative Abundance	Oligohaline Indicative Abundance	Oligohaline Sensitive Abundance	Tanypodinae to Chironomidae Ratio	Abundance Deep Deposit Feeders	Abundance Carnivore/Omnivores
Potomac River									
36	0.00	-42.27	0.004	0.56	NA	NA	NA	0.50	NA
40	0.00	-1.97	0.00	NA	-0.82	0.00	0.00	NA	0.59
Patuxent River									
79	0.00	235.99	-0.005	-1.90	NA	NA	NA	-0.33	NA
Choptank River									
66	0.00	92.12	0.12	NA	0.56	0.00	0.00	NA	1.09
Maryland Western Shore Tributaries									
203(a)	0.02	109.10	-0.001	NA	-0.01	0.00	0.22	NA	0.56
Maryland Eastern Shore Tributaries									
29	0.02	-65.56	-0.10	NA	-3.35	0.15	0.00	NA	0.17

APPENDIX B

FIXED SITE B-IBI VALUES, SUMMER 2003

Appendix Table B-1. Fixed site B-IBI values, Summer 2003

Station	Sampling Date	Latitude (NAD83 Decimal Degrees)	Longitude (NAD83 Decimal Degrees)	B-IBI	Status
001	28-Aug-03	38.41918	-76.4182	2.56	Degraded
006	28-Aug-03	38.44223	-76.4442	2.33	Degraded
015	28-Aug-03	38.71427	-76.5127	3.56	Meets Goal
022	2-Sep-03	39.25450	-76.5877	1.00	Severely Degraded
023	2-Sep-03	39.20822	-76.5237	2.33	Degraded
024	2-Sep-03	39.12183	-76.3556	2.56	Degraded
026	3-Sep-03	39.27142	-76.2901	3.93	Meets Goal
029	3-Sep-03	39.47953	-75.9449	2.56	Degraded
036	11-Sep-03	38.77000	-77.0371	1.50	Severely Degraded
040	26-Aug-03	38.35733	-77.2308	3.49	Meets Goal
043	26-Aug-03	38.38430	-76.9887	3.27	Meets Goal
044	26-Aug-03	38.38550	-76.9960	1.80	Severely Degraded
047	26-Aug-03	38.36508	-76.9841	3.53	Meets Goal
051	26-Aug-03	38.20533	-76.7383	2.78	Marginal
052	26-Aug-03	38.19217	-76.7480	1.44	Severely Degraded
062	16-Sep-03	38.38412	-75.8506	3.80	Meets Goal
064	28-Aug-03	38.59033	-76.0697	2.44	Degraded
066	10-Sep-03	38.80098	-75.9220	3.22	Meets Goal
068	10-Sep-03	39.13252	-76.0776	3.13	Meets Goal
071	29-Aug-03	38.39500	-76.5492	1.89	Severely Degraded
074	29-Aug-03	38.54883	-76.6765	3.00	Meets Goal
077	29-Aug-03	38.60433	-76.6753	4.07	Meets Goal
079	11-Sep-03	38.74925	-76.6903	1.67	Severely Degraded
201	2-Sep-03	39.23418	-76.4975	1.40	Severely Degraded
202	2-Sep-03	39.21763	-76.5642	1.27	Severely Degraded
203	8-Sep-03	39.27500	-76.4444	2.67	Marginal
204	4-Sep-03	39.00670	-76.5050	2.11	Degraded

APPENDIX C

RANDOM SITE B-IBI VALUES, SUMMER 2003

Appendix Table C-1. Random site B-IBI values, Summer 2003					
Station	Sampling Date	Latitude (NAD83 Decimal Degrees)	Longitude (NAD83 Decimal Degrees)	B-IBI	Status
MET-10401	16-Sep-03	37.97391667	-75.64466667	2.00	Severely Degraded
MET-10402	25-Aug-03	38.03413333	-75.84383333	3.33	Meets Goal
MET-10403	25-Aug-03	38.039998	-75.84621	4.00	Meets Goal
MET-10405	25-Aug-03	38.120816	-75.907811	3.33	Meets Goal
MET-10406	25-Aug-03	38.121987	-75.85571	2.33	Degraded
MET-10407	25-Aug-03	38.12598333	-75.86061667	3.33	Meets Goal
MET-10408	25-Aug-03	38.12898333	-75.86443333	3.33	Meets Goal
MET-10409	25-Aug-03	38.21993333	-75.86696667	1.80	Severely Degraded
MET-10410	25-Aug-03	38.231305	-75.868991	2.60	Degraded
MET-10411	16-Sep-03	38.32403333	-75.91958333	2.60	Degraded
MET-10412	16-Sep-03	38.3589	-75.86176667	2.60	Degraded
MET-10413	28-Aug-03	38.591665	-76.111311	3.00	Meets Goal
MET-10414	28-Aug-03	38.625632	-76.147986	2.20	Degraded
MET-10415	28-Aug-03	38.62573333	-76.12113333	2.60	Degraded
MET-10416	10-Sep-03	38.77573333	-75.9679	2.50	Degraded
MET-10417	3-Sep-03	38.99505	-76.1936	1.00	Severely Degraded
MET-10418	3-Sep-03	38.99858333	-76.22785	1.67	Severely Degraded
MET-10419	3-Sep-03	39.00116667	-76.23431667	1.67	Severely Degraded
MET-10420	3-Sep-03	39.0135	-76.17898333	1.40	Severely Degraded
MET-10421	3-Sep-03	39.02155	-76.24965	2.60	Degraded
MET-10422	3-Sep-03	39.08481667	-76.20036667	2.20	Degraded
MET-10423	10-Sep-03	39.2257	-76.03011667	3.33	Meets Goal
MET-10424	3-Sep-03	39.48513333	-75.9367	3.00	Meets Goal
MET-10425	3-Sep-03	39.50825	-75.89733333	4.67	Meets Goal
MET-10426	3-Sep-03	39.02785	-76.18083333	2.20	Degraded
MMS-10501	16-Sep-03	37.93031667	-75.81063333	3.00	Meets Goal
MMS-10503	25-Aug-03	37.964241	-76.084982	2.67	Marginal
MMS-10504	25-Aug-03	38.01041667	-75.8854	3.67	Meets Goal
MMS-10505	15-Sep-03	38.009102	-75.880892	2.00	Severely Degraded
MMS-10506	25-Aug-03	38.021389	-76.148557	1.33	Severely Degraded
MMS-10507	25-Aug-03	38.037618	-76.083115	2.60	Degraded
MMS-10508	25-Aug-03	38.055238	-76.137945	2.33	Degraded
MMS-10509	25-Aug-03	38.06505	-75.88678333	3.33	Meets Goal
MMS-10510	25-Aug-03	38.075408	-75.936214	3.00	Meets Goal
MMS-10511	15-Sep-03	38.094419	-76.181967	3.00	Meets Goal
MMS-10512	15-Sep-03	38.094419	-76.161921	3.00	Meets Goal
MMS-10513	15-Sep-03	38.138352	-76.1631	1.67	Severely Degraded
MMS-10514	25-Aug-03	38.19096667	-75.94928333	3.33	Meets Goal
MMS-10515	25-Aug-03	38.205817	-76.09019	3.00	Meets Goal
MMS-10517	25-Aug-03	38.22401667	-76.14496667	2.60	Degraded

Appendix Table C-1. (Continued)

Station	Sampling Date	Latitude (NAD83 Decimal Degrees)	Longitude (NAD83 Decimal Degrees)	B-IBI	Status
MMS-10518	25-Aug-03	38.25408333	-76.10895	2.20	Degraded
MMS-10519	28-Aug-03	38.553692	-76.407871	2.67	Marginal
MMS-10520	28-Aug-03	38.634719	-76.337515	1.00	Severely Degraded
MMS-10521	28-Aug-03	38.64237	-76.16821	2.60	Degraded
MMS-10522	28-Aug-03	38.774634	-76.512028	1.00	Severely Degraded
MMS-10523	09-Sep-03	38.7906	-76.2077	1.00	Severely Degraded
MMS-10524	04-Sep-03	38.85398333	-76.4845	2.20	Degraded
MMS-10525	04-Sep-03	38.87553333	-76.37568333	3.40	Meets Goal
MMS-10526	28-Aug-03	38.720036	-76.272269	3.00	Meets Goal
MMS-10527	25-Aug-03	38.052456	-76.1292	3.00	Meets Goal
MWT-10301	04-Sep-03	38.86833333	-76.50146667	3.40	Meets Goal
MWT-10302	04-Sep-03	38.87611667	-76.52358333	2.20	Degraded
MWT-10303	04-Sep-03	38.87743333	-76.48933333	2.60	Degraded
MWT-10304	04-Sep-03	38.90771667	-76.48903333	2.60	Degraded
MWT-10305	04-Sep-03	38.93873333	-76.53405	2.20	Degraded
MWT-10306	04-Sep-03	38.95016667	-76.53998333	3.40	Meets Goal
MWT-10307	04-Sep-03	38.9537	-76.55696667	1.80	Severely Degraded
MWT-10308	04-Sep-03	38.98783333	-76.48396667	2.20	Degraded
MWT-10309	04-Sep-03	39.04576667	-76.53835	2.20	Degraded
MWT-10310	04-Sep-03	39.07028333	-76.46636667	2.20	Degraded
MWT-10311	04-Sep-03	39.07686667	-76.49973333	1.00	Severely Degraded
MWT-10312	02-Sep-03	39.16233333	-76.45817	3.00	Meets Goal
MWT-10313	02-Sep-03	39.18518333	-76.46965	3.00	Meets Goal
MWT-10314	02-Sep-03	39.1855	-76.51465	4.20	Meets Goal
MWT-10315	02-Sep-03	39.1905	-76.46448333	3.00	Meets Goal
MWT-10316	02-Sep-03	39.19225	-76.5044	3.40	Meets Goal
MWT-10317	02-Sep-03	39.196	-76.57445	1.80	Severely Degraded
MWT-10318	02-Sep-03	39.21475	-76.52503333	1.00	Severely Degraded
MWT-10319	08-Sep-03	39.24866667	-76.41993333	2.67	Marginal
MWT-10320	08-Sep-03	39.31296667	-76.30218333	3.33	Meets Goal
MWT-10321	08-Sep-03	39.33663333	-76.30118333	4.00	Meets Goal
MWT-10324	08-Sep-03	39.38216667	-76.32628333	2.50	Degraded
MWT-10326	02-Sep-03	39.21426667	-76.54518333	2.60	Degraded
MWT-10329	08-Sep-03	39.32498333	-76.31558333	2.67	Marginal
MWT-10330	08-Sep-03	39.33986667	-76.30338333	3.33	Meets Goal
PMR-10101	15-Sep-03	38.017047	-76.334075	1.00	Severely Degraded
PMR-10102	15-Sep-03	38.025951	-76.469186	1.40	Severely Degraded
PMR-10103	15-Sep-03	38.042069	-76.380545	1.00	Severely Degraded
PMR-10104	27-Aug-03	38.072727	-76.467991	1.00	Severely Degraded
PMR-10105	27-Aug-03	38.091212	-76.466677	1.33	Severely Degraded

Appendix Table C-1. (Continued)

Station	Sampling Date	Latitude (NAD83 Decimal Degrees)	Longitude (NAD83 Decimal Degrees)	B-IBI	Status
PMR-10106	27-Aug-03	38.117812	-76.443502	1.00	Severely Degraded
PMR-10107	27-Aug-03	38.13048333	-76.49058333	2.60	Degraded
PMR-10108	26-Aug-03	38.164475	-76.736421	1.00	Severely Degraded
PMR-10109	27-Aug-03	38.167405	-76.576701	1.00	Severely Degraded
PMR-10110	15-Sep-03	38.171125	-76.602027	1.00	Severely Degraded
PMR-10111	26-Aug-03	38.190286	-76.792568	1.00	Severely Degraded
PMR-10112	15-Sep-03	38.192991	-76.613615	1.00	Severely Degraded
PMR-10113	27-Aug-03	38.196034	-76.442307	1.00	Severely Degraded
PMR-10114	26-Aug-03	38.208207	-76.870935	1.00	Severely Degraded
PMR-10115	26-Aug-03	38.217112	-76.809771	1.00	Severely Degraded
PMR-10116	26-Aug-03	38.281245	-76.957903	1.80	Severely Degraded
PMR-10117	26-Aug-03	38.283837	-76.835216	1.40	Severely Degraded
PMR-10118	26-Aug-03	38.320018	-76.998639	2.60	Degraded
PMR-10119	26-Aug-03	38.339292	-77.228961	3.33	Meets Goal
PMR-10120	26-Aug-03	38.347858	-77.177951	3.00	Meets Goal
PMR-10121	26-Aug-03	38.366794	-77.286541	4.00	Meets Goal
PMR-10122	26-Aug-03	38.38478333	-77.14673333	4.00	Meets Goal
PMR-10123	26-Aug-03	38.387307	-77.256318	5.00	Meets Goal
PMR-10124	26-Aug-03	38.403989	-77.02516	2.20	Degraded
PMR-10125	26-Aug-03	38.447045	-77.026235	2.00	Severely Degraded
PXR-10201	28-Aug-03	38.295366	-76.448438	2.20	Degraded
PXR-10203	28-Aug-03	38.317667	-76.435041	2.20	Degraded
PXR-10204	28-Aug-03	38.319525	-76.426483	2.60	Degraded
PXR-10205	29-Aug-03	38.339348	-76.49495	3.00	Meets Goal
PXR-10206	29-Aug-03	38.35016667	-76.48245	3.00	Meets Goal
PXR-10207	29-Aug-03	38.361287	-76.49574	1.00	Severely Degraded
PXR-10208	29-Aug-03	38.365675	-76.505746	1.80	Severely Degraded
PXR-10209	29-Aug-03	38.370888	-76.517366	1.80	Severely Degraded
PXR-10210	29-Aug-03	38.372385	-76.511276	1.80	Severely Degraded
PXR-10211	29-Aug-03	38.388698	-76.504496	2.60	Degraded
PXR-10212	29-Aug-03	38.403255	-76.589223	2.60	Degraded
PXR-10213	29-Aug-03	38.407746	-76.5779	1.80	Severely Degraded
PXR-10214	29-Aug-03	38.40914	-76.587709	1.00	Severely Degraded
PXR-10215	29-Aug-03	38.414663	-76.580632	1.40	Severely Degraded
PXR-10216	29-Aug-03	38.417296	-76.583594	1.80	Severely Degraded
PXR-10217	29-Aug-03	38.41881667	-76.56738333	3.40	Meets Goal
PXR-10219	29-Aug-03	38.443933	-76.626518	1.00	Severely Degraded
PXR-10220	29-Aug-03	38.44463333	-76.60243333	2.60	Degraded
PXR-10221	29-Aug-03	38.454618	-76.620198	2.60	Degraded
PXR-10223	29-Aug-03	38.53285	-76.67493333	3.40	Meets Goal

Appendix Table C-1. (Continued)					
Station	Sampling Date	Latitude (NAD83 Decimal Degrees)	Longitude (NAD83 Decimal Degrees)	B-IBI	Status
PXR-10224	29-Aug-03	38.627137	-76.679909	3.00	Meets Goal
PXR-10225	11-Sep-03	38.76161667	-76.69908333	2.50	Degraded
PXR-10226	29-Aug-03	38.413786	-76.584022	1.00	Severely Degraded
PXR-10227	11-Sep-03	38.71871667	-76.69823333	3.50	Meets Goal
PXR-10228	29-Aug-03	38.37206667	-76.48991667	2.20	Degraded
UPB-10601	03-Sep-03	39.06123333	-76.2486	2.60	Degraded
UPB-10602	02-Sep-03	39.07483333	-76.31298333	1.67	Severely Degraded
UPB-10603	02-Sep-03	39.07983333	-76.37995	3.00	Meets Goal
UPB-10604	03-Sep-03	39.08475	-76.26416667	1.80	Severely Degraded
UPB-10605	02-Sep-03	39.08928333	-76.34591667	2.60	Degraded
UPB-10606	02-Sep-03	39.08933333	-76.33443333	1.80	Severely Degraded
UPB-10607	02-Sep-03	39.09133333	-76.3644	2.60	Degraded
UPB-10608	02-Sep-03	39.12055	-76.33753333	3.00	Meets Goal
UPB-10609	02-Sep-03	39.12745	-76.4169	3.00	Meets Goal
UPB-10610	02-Sep-03	39.14511667	-76.34526667	3.40	Meets Goal
UPB-10611	02-Sep-03	39.16936667	-76.41828333	3.00	Meets Goal
UPB-10612	02-Sep-03	39.17916667	-76.44035	1.67	Severely Degraded
UPB-10613	02-Sep-03	39.20983333	-76.35728333	2.20	Degraded
UPB-10615	03-Sep-03	39.23435	-76.2465	4.60	Meets Goal
UPB-10616	03-Sep-03	39.29438333	-76.18886667	3.00	Meets Goal
UPB-10617	03-Sep-03	39.2947	-76.24708333	3.80	Meets Goal
UPB-10618	08-Sep-03	39.30823333	-76.35583333	4.00	Meets Goal
UPB-10620	03-Sep-03	39.3854	-76.14435	5.00	Meets Goal
UPB-10621	03-Sep-03	39.47885	-75.99811667	4.00	Meets Goal
UPB-10622	03-Sep-03	39.48316667	-76.09066667	3.50	Meets Goal
UPB-10623	03-Sep-03	39.47351667	-76.01706667	3.50	Meets Goal
UPB-10624	03-Sep-03	39.51845	-75.9878	2.00	Severely Degraded
UPB-10625	03-Sep-03	39.53778333	-76.0629	3.00	Meets Goal
UPB-10626	02-Sep-03	39.06301667	-76.37138333	1.80	Severely Degraded
UPB-10627	02-Sep-03	39.11256667	-76.34516667	2.60	Degraded